

MIS12 (FL-205): sc-98368

BACKGROUND

Chromosome segregation requires assembly of kinetochores on centromeric chromatin to mediate interactions with spindle microtubules and control cell-cycle progression. MIS12 (minichromosome instability 12), also known as MTW1, hMis12, KNTC2AP or MIND kinetochore complex component homolog, is a 205 amino acid nuclear protein that is associated with the kinetochore. MIS12 is a component of the MIS12 complex, which is required for kinetochore formation during mitosis and normal chromosome alignment and segregation. The MIS12 complex consists of MIS12, DSN1, NSL1 and PMF-1. MIS12 is part of a network of complexes that provide microtubule attachment and generates pulling forces from depolymerization. MIS12 is encoded by a gene located on human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

REFERENCES

- Goshima, G., et al. 2003. Human centromere chromatin protein hMIS12, essential for equal segregation, is independent of CENP-A loading pathway. *J. Cell Biol.* 160: 25-39.
- Obuse, C., et al. 2004. A conserved MIS12 centromere complex is linked to heterochromatic HP1 and outer kinetochore protein ZWINT-1. *Nat. Cell Biol.* 6: 1135-1141.
- Kline, S.L., et al. 2006. The human MIS12 complex is required for kinetochore assembly and proper chromosome segregation. *J. Cell Biol.* 173: 9-17.
- Liu, S.T., et al. 2006. Mapping the assembly pathways that specify formation of the trilaminar kinetochore plates in human cells. *J. Cell Biol.* 175: 41-53.
- Zhang, R., et al. 2007. HP1 proteins are essential for a dynamic nuclear response that rescues the function of perturbed heterochromatin in primary human cells. *Mol. Cell Biol.* 27: 949-962.
- Hemmerich, P., et al. 2008. Dynamics of inner kinetochore assembly and maintenance in living cells. *J. Cell Biol.* 180: 1101-1114.
- Wan, X., et al. 2009. Protein architecture of the human kinetochore microtubule attachment site. *Cell* 137: 672-684.
- Li, X. and Dawe, R.K. 2009. Fused sister kinetochores initiate the reductional division in meiosis I. *Nat. Cell Biol.* 11: 1103-1108.

CHROMOSOMAL LOCATION

Genetic locus: MIS12 (human) mapping to 17p13.2; Mis12 (mouse) mapping to 11 B4.

SOURCE

MIS12 (FL-205) is a rabbit polyclonal antibody raised against amino acids 1-205 representing full length MIS12 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

MIS12 (FL-205) is recommended for detection of MIS12 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MIS12 (FL-205) is also recommended for detection of MIS12 in additional species, including equine and bovine.

Suitable for use as control antibody for MIS12 siRNA (h): sc-93889, MIS12 siRNA (m): sc-149443, MIS12 shRNA Plasmid (h): sc-93889-SH, MIS12 shRNA Plasmid (m): sc-149443-SH, MIS12 shRNA (h) Lentiviral Particles: sc-93889-V and MIS12 shRNA (m) Lentiviral Particles: sc-149443-V.


Molecular Weight of MIS12: 25 kDa.

Positive Controls: rat small intestine extract: sc-364811.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MIS12 (FL-205): sc-98368. Western blot analysis of MIS12 expression in rat small intestine tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.