

# Otoferlin (H-150): sc-98457

## BACKGROUND

Otoferlin is a single-pass type II membrane protein composed of 1,230 amino acid residues. Otoferlin exists in four isoforms; isoform 1 is the full-length Otoferlin protein, whereas isoforms 2-4 are shorter versions of the protein. Expression of isoforms 1 and 3 is demonstrated in adult brain, while isoform 2 is expressed in the fetus, adult brain, heart, placenta, skeletal muscle and kidney tissues. Otoferlin has three C2 domains and a single carboxy-terminal transmembrane domain. The Otoferlin gene, OTOF, and its surrounding genes map to 2p23.3, and the 5' region of OTOF is centromeric. Mutations in the OTOF gene are implicated in deafness. Otoferlin is homologous to the *C. elegans* spermatogenesis factor FER-1 and to human dysferlin, implicating the involvement of Otoferlin in the Ca<sup>2+</sup>-triggered synaptic vesicle-plasma membrane fusion.

## REFERENCES

1. Starr, A., Sininger, Y., Winter, M., Derebery, M.J., Oba, S. and Michalewski, H.J. 1998. Transient deafness due to temperature-sensitive auditory neuropathy. *Ear Hear.* 19: 169-179.
2. Yasunaga, S., Grati, M., Cohen-Salmon, M., El-Amraoui, A., Mustapha, M., Salem, N., El-Zir, E., Loiselet, J. and Petit, C. 1999. A mutation in OTOF, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic form of deafness. *Nat. Genet.* 21: 363-369.
3. Adato, A., Raskin, L., Petit, C. and Bonne-Tamir, B. 2000. Deafness heterogeneity in a Druze isolate from the Middle East: novel OTOF and PDS mutations, low prevalence of GJB2 35delG mutation and indication for a new DFNB locus. *Eur. J. Hum. Genet.* 8: 437-442.
4. Migliosi, V., Modamio-Høybjør, S., Moreno-Pelayo, M.A., Rodríguez-Ballesteros, M., Villamar, M., Tellería, D., Menendez, I., Moreno, F. and Del Castillo, I. 2002. Q829X, a novel mutation in the gene encoding otoferlin (OTOF), is frequently found in Spanish patients with prelingual non-syndromic hearing loss. *J. Med. Genet.* 39: 502-506.
5. Mirghomizadeh, F., Pfister, M., Apaydin, F., Petit, C., Kupka, S., Pusch, C.M., Zenner, H.P. and Blin, N. 2002. Substitutions in the conserved C2C domain of OTOF of nonsyndromic autosomal recessive deafness. *Neurobiol. Dis.* 10: 157-164.
6. Mirghomizadeh, F., Pfister, M., Blin, N. and Pusch, C.M. 2002. Uncommon cytidine-homopolymer dimorphism in 5'-UTR of the human otoferlin gene. *Int. J. Mol. Med.* 11: 63-64.
7. Rodríguez-Ballesteros, M., del Castillo, F.J., Martín, Y., Moreno-Pelayo, M.A., Morera, C., Prieto, F., Marco, J., Morant, A., Gallo-Terán, J., Morales-Angulo, C., Navas, C., Trinidad, G., Tapia, M.C., Moreno, F. and del Castillo, I. 2003. Auditory neuropathy in patients carrying mutations in the otoferlin gene (OTOF). *Hum. Mutat.* 22: 451-456.

## CHROMOSOMAL LOCATION

Genetic locus: OTOF (human) mapping to 2p23.3; Otof (mouse) mapping to 5 B1.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## SOURCE

Otoferlin (H-150) is a rabbit polyclonal antibody raised against amino acids 691-840 mapping within an internal region of Otoferlin of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Otoferlin (H-150) is recommended for detection of Otoferlin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Otoferlin (H-150) is also recommended for detection of Otoferlin in additional species, including equine, canine and bovine.

Suitable for use as control antibody for Otoferlin siRNA (h): sc-61269, Otoferlin siRNA (m): sc-61270, Otoferlin shRNA Plasmid (h): sc-61269-SH, Otoferlin shRNA Plasmid (m): sc-61270-SH, Otoferlin shRNA (h) Lentiviral Particles: sc-61269-V and Otoferlin shRNA (m) Lentiviral Particles: sc-61270-V.

Molecular Weight of Otoferlin: 140.5 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409 or DU 145 cell lysate: sc-2268.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **Otoferlin (C-12): sc-271092**, our highly recommended monoclonal alternative to Otoferlin (H-150).