

## TRF4 (H-172): sc-98490

### BACKGROUND

DNA replication, recombination and repair, all of which are necessary for genomic stability, require the presence of exonucleases. In DNA replication, these enzymes are involved in the processing of Okazaki fragments, whereas in DNA repair, they function to excise damaged DNA fragments and correct recombinational mismatches. These exonucleases include the family of DNA polymerases. TRF4 (topoisomerase-related function protein 4-1), also known as POLS (polymerase  $\sigma$ ), POLK (polymerase  $\kappa$ ) or LAK-1, is a member of the Y-family of translesion DNA polymerases and is involved in DNA replication and repair. In addition, TRF4 may be involved in the cohesion of sister chromatids in association with the replication fork. Once replicated, sister chromatids must be held together until anaphase to ensure accurate chromosome segregation.

### REFERENCES

1. Walowsky, C., Fitzhugh, D.J., Castaño, I.B., Ju, J.Y., Levin, N.A. and Christman, M.F. 1999. The topoisomerase-related function gene TRF4 affects cellular sensitivity to the antitumor agent camptothecin. *J. Biol. Chem.* 274: 7302-7308.
2. Wang, Z., Castaño, I.B., De Las Peñas, A., Adams, C. and Christman, M.F. 2000. Pol  $\kappa$ : a DNA polymerase required for sister chromatid cohesion. *Science* 289: 774-779.
3. Carson, D.R. and Christman, M.F. 2001. Evidence that replication fork components catalyze establishment of cohesion between sister chromatids. *Proc. Natl. Acad. Sci. USA* 98: 8270-8275.
4. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605198. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Hubscher, U., Maga, G. and Spadari, S. 2002. Eukaryotic DNA polymerases. *Annu. Rev. Biochem.* 71: 133-163.
6. Wang, Z. and Christman, M.F. 2002. Replication-related activities establish cohesion between sister chromatids. *Cell Biochem. Biophys.* 35: 289-301.
7. Choi, J.Y., Angel, K.C. and Guengerich, F.P. 2006. Translesion synthesis across bulky N2-alkyl guanine DNA adducts by human DNA polymerase  $\kappa$ . *J. Biol. Chem.* 281: 21062-21072.
8. Yasui, M., Suzuki, N., Liu, X., Okamoto, Y., Kim, S.Y., Laxmi, Y.R. and Shibutani, S. 2007. Mechanism of translesion synthesis past an equine estrogen-DNA adduct by Y-family DNA polymerases. *J. Mol. Biol.* 371: 1151-1162.

### CHROMOSOMAL LOCATION

Genetic locus: POLS (human) mapping to 5p15.31; Pols (mouse) mapping to 13 C1.

### SOURCE

TRF4 (H-172) is a rabbit polyclonal antibody raised against amino acids 371-542 mapping at the C-terminus of TRF4 of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### APPLICATIONS

TRF4 (H-172) is recommended for detection of TRF4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TRF4 (H-172) is also recommended for detection of TRF4 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for TRF4 siRNA (h): sc-91936, TRF4 siRNA (m): sc-154635, TRF4 shRNA Plasmid (h): sc-91936-SH, TRF4 shRNA Plasmid (m): sc-154635-SH, TRF4 shRNA (h) Lentiviral Particles: sc-91936-V and TRF4 shRNA (m) Lentiviral Particles: sc-154635-V.

Molecular Weight of TRF4: 60 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker<sup>™</sup> compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz<sup>™</sup> Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **TRF4 (2794C4a): sc-81637**, our highly recommended monoclonal alternative to TRF4 (H-172).