



CLEC-2B (H-50): sc-98521

BACKGROUND

The C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily consists of a variety of proteins that share a common protein fold and have diverse functions, including cell-cell signaling, cell adhesion, glycoprotein turnover and immune responses. CLEC-2B (C-type lectin domain family 2 member B), also known as IFNRG1 (IFN- α 2b-inducing-related protein 1) and AICL (Activation-induced C-type lectin), is a 149 amino acid membrane protein that contains one extracellular C-type lectin domain near its carboxyl-terminus. The CLEC-2B gene is located within the natural killer gene complex, which includes genes encoding for CD69, CLEC-2D, CD161 and CD94. Since downregulation of CLEC-2B reduces NK cell-mediated cytotoxicity towards target cells and it is expressed in lymphoid tissues, CLEC-2B is thought play a role in the immune system.

REFERENCES

1. Hamann, J., Montgomery, K.T., Lau, S., Kucherlapati, R. and van Lier, R.A. 1997. AICL: a new activation-induced antigen encoded by the human NK gene complex. *Immunogenetics* 45: 295-300.
2. Yokoyama-Kobayashi, M., Yamaguchi, T., Sekine, S. and Kato, S. 1999. Selection of cDNAs encoding putative type II membrane proteins on the cell surface from a human full-length cDNA bank. *Gene* 228: 161-167.
3. Renedo, M., Arce, I., Montgomery, K., Roda-Navarro, P., Lee, E., Kucherlapati, R. and Fernández-Ruiz, E. 2000. A sequence-ready physical map of the region containing the human natural killer gene complex on chromosome 12p12.3-p13.2. *Genomics* 65: 129-136.
4. Eichler, W., Ruschpler, P., Wobus, M. and Drössler, K. 2001. Differentially induced expression of C-type lectins in activated lymphocytes. *J. Cell. Biochem. Suppl.* 36: 201-208.
5. Welte, S., Kuttruff, S., Waldhauer, I. and Steinle, A. 2006. Mutual activation of natural killer cells and monocytes mediated by NKp80-AICL interaction. *Nat. Immunol.* 7: 1334-1342.
6. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 603242. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Thomas, M., Wills, M. and Lehner, P.J. 2008. Natural killer cell evasion by an E3 ubiquitin ligase from Kaposi's sarcoma-associated herpesvirus. *Biochem. Soc. Trans.* 36: 459-463.
8. Kuttruff, S., Koch, S., Kelp, A., Pawelec, G., Rammensee, H.G. and Steinle, A. 2009. NKp80 defines and stimulates a reactive subset of CD8 T cells. *Blood* 113: 358-369.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: CLEC2B (human) mapping to 12p13.31.

SOURCE

CLEC-2B (H-50) is a rabbit polyclonal antibody raised against amino acids 41-90 mapping within an internal region of CLEC-2B of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CLEC-2B (H-50) is recommended for detection of CLEC-2B of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CLEC-2B siRNA (h): sc-95808, CLEC-2B shRNA Plasmid (h): sc-95808-SH and CLEC-2B shRNA (h) Lentiviral Particles: sc-95808-V.

Molecular Weight of CLEC-2B: 17 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.