

TFIIIB90-1/2/3/5 (H-40): sc-98583

BACKGROUND

RNA polymerase (pol) III synthesizes tRNA, 5s rRNA, 7SL RNA and U6 snRNA and is overexpressed in many transformed cell lines and tumors *in vivo*, since cells must duplicate its protein components before division. Therefore, in order to maintain rapid growth, cells must produce a high level of Pol III transcribed RNA, which requires the presence of the TFIIIB and TFIIIC2 transcription factor complexes. The TFIIIC2 complex is composed of five subunits, TFIIIC220, TFIIIC110, TFIIIC102, TFIIIC90 and TFIIIC63, that are overexpressed in adenovirus transformed cells as well as in malignant cells *in vivo*, such as ovarian carcinomas. TFIIIC2 recruits RNA pol III and TFIIIB to promoter elements and may be a key component in the deregulation of malignant cells. The TFIIIB complex includes the TATA-binding protein (TBP), TFIIIB-related factor 1 (TFIIIB90, BRF1) and TFIIIB, the expression of which are also upregulated in transformed cells. In many carcinomas, the tumor suppressors retinoblastoma (RB) and p53 are inactivated, which affects their ability to bind and inactivate the function of TFIIIB.

REFERENCES

1. Scott, M.R., et al. 1983. Activation of mouse genes in transformed cells. *Cell* 34: 557-567.
2. Chen, W., et al. 1997. Expression of neural BC1 RNA: induction in murine tumours. *Eur. J. Cancer* 33: 288-292.
3. Hsieh, Y.J., et al. 1999. The TFIIIC90 subunit of TFIIIC interacts with multiple components of the RNA polymerase III machinery and contains a histone-specific acetyltransferase activity. *Mol. Cell. Biol.* 19: 7697-7704.
4. Winter, A.G., et al. 2000. RNA polymerase III transcription factor TFIIIC2 is overexpressed in ovarian tumors. *Proc. Natl. Acad. Sci. USA* 97: 12619-12624.
5. Moir, R.D., et al. 2000. Interactions between the tetratricopeptide repeat-containing transcription factor TFIIIC131 and its ligand, TFIIIB70. Evidence for a conformational change in the complex. *J. Biol. Chem.* 275: 26591-26598.

CHROMOSOMAL LOCATION

Genetic locus: BRF1 (human) mapping to 14q32.33; Brf1 (mouse) mapping to 12 F1.

SOURCE

TFIIIB90-1/2/3/5 (H-40) is a rabbit polyclonal antibody raised against amino acids 251-290 mapping within an internal region of TFIIIB90 -1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-98583 X, 200 µg/0.1 ml.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

TFIIIB90-1/2/3/5 (H-40) is recommended for detection of TFIIIB90 isoforms 1, 2, 3 and 5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TFIIIB90-1/2/3/5 (H-40) is also recommended for detection of TFIIIB90 isoforms 1, 2, 3 and 5 in additional species, including equine, canine, bovine and avian.

Suitable for use as control antibody for TFIIIB90 siRNA (h): sc-38535, TFIIIB90 siRNA (m): sc-154232, TFIIIB90 shRNA Plasmid (h): sc-38535-SH, TFIIIB90 shRNA Plasmid (m): sc-154232-SH, TFIIIB90 shRNA (h) Lentiviral Particles: sc-38535-V and TFIIIB90 shRNA (m) Lentiviral Particles: sc-154232-V.

TFIIIB90-1/2/3/5 (H-40) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TFIIIB90-1/2/3/5: 74 kDa.

Positive Controls: TFIIIB90-1 (h): 293T Lysate: sc-111635 or Hep G2 nuclear extract: sc-364819.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TFIIIB90-1/2/3/5 (H-40): sc-98583. Western blot analysis of TFIIIB90-1 expression in non-transfected: sc-117752 (A) and human TFIIIB90-1 transfected: sc-111635 (B) 293T whole cell lysates and Hep G2 nuclear extract (C).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.