

Rac GAP1 (H-300): sc-98617

BACKGROUND

A large number of low molecular weight, GTP binding proteins of the Ras superfamily have been identified. These proteins regulate many fundamental processes in all eukaryotic cells such as growth, vesicle traffic and cytoskeletal organization. GTPase-activating proteins (GAPs) accelerate the intrinsic rate of GTP hydrolysis of Ras-related proteins, resulting in downregulation of their active form. Through this function, GAPs negatively regulate Ras-mediated signaling. Rac GAP1 (Rac GTPase activating protein 1), also known as MgcRacGAP (male germ cell Rac GTPase activating protein), ID-GAP or HsCYK-4, functions as a GAP and exhibits strong activity towards Rac 1 and Cdc42. Highly expressed in thymus, placenta and testis with lower levels in spleen and peripheral blood lymphocytes, Rac GAP1 contains one Rho GAP domain and one phorbol-ester/DAG-type zinc finger. Rac GAP1 plays an essential role in cytokinesis, functioning as a scaffold protein as well as a GTPase regulator. During cytokinesis, Rac GAP1 is phosphorylated at multiple sites.

CHROMOSOMAL LOCATION

Genetic locus: RACGAP1 (human) mapping to 12q13.12; Racgap1 (mouse) mapping to 15 F1.

SOURCE

Rac GAP1 (H-300) is a rabbit polyclonal antibody raised against amino acids 142-437 mapping within an internal region of Rac GAP1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Rac GAP1 (H-300) is recommended for detection of Rac GAP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Rac GAP1 (H-300) is also recommended for detection of Rac GAP1 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for Rac GAP1 siRNA (h): sc-76335, Rac GAP1 siRNA (m): sc-76336, Rac GAP1 shRNA Plasmid (h): sc-76335-SH, Rac GAP1 shRNA Plasmid (m): sc-76336-SH, Rac GAP1 shRNA (h) Lentiviral Particles: sc-76335-V and Rac GAP1 shRNA (m) Lentiviral Particles: sc-76336-V.


Molecular Weight of Rac GAP1: 70 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Rac GAP1 (H-300): sc-98617. Western blot analysis of Rac GAP1 expression in Jurkat (A) and K-562 (B) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **Rac GAP1 (A-6): sc-271110** or **Rac GAP1 (B-7): sc-166477**, our highly recommended monoclonal alternatives to Rac GAP1 (H-300).