

ALP (H-157): sc-98652

BACKGROUND

ALP (actinin-associated LIM protein), also known as PDLIM3 (PDZ and LIM domain protein 3) is a 364 amino acid protein that contains one LIM zinc-binding domain and one PDZ domain and localizes to the cytoplasm, as well as to myofiber Z-lines. Existing as three alternatively spliced isoforms, two of which exhibit tissue-specific expression in skeletal muscle and heart, ALP interacts with α -actinin-2 and, via this interaction, is thought to play a role in actin filament organization, specifically regulating the association of actin filaments arrays with muscle cells. The gene encoding ALP maps to a region on human chromosome 4 that is associated with facioscapulohumeral muscular dystrophy, suggesting that defects in the ALP gene may be involved in the pathogenesis of muscular dystrophy.

REFERENCES

1. Piétu, G., et al. 1996. Novel gene transcripts preferentially expressed in human muscles revealed by quantitative hybridization of a high density cDNA array. *Genome Res.* 6: 492-503.
2. Xia, H., et al. 1997. Actinin-associated LIM protein: identification of a domain interaction between PDZ and spectrin-like repeat motifs. *J. Cell Biol.* 139: 507-515.

CHROMOSOMAL LOCATION

Genetic locus: PDLIM3 (human) mapping to 4q35.1; Pdlim3 (mouse) mapping to 8 B1.1.

SOURCE

ALP (H-157) is a rabbit polyclonal antibody raised against amino acids 44-200 mapping within an internal region of ALP of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

ALP (H-157) is recommended for detection of ALP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ALP (H-157) is also recommended for detection of ALP in additional species, including equine, canine and porcine.

Suitable for use as control antibody for ALP siRNA (h): sc-72485, ALP siRNA (m): sc-72486, ALP shRNA Plasmid (h): sc-72485-SH, ALP shRNA Plasmid (m): sc-72486-SH, ALP shRNA (h) Lentiviral Particles: sc-72485-V and ALP shRNA (m) Lentiviral Particles: sc-72486-V.


Molecular Weight of ALP: 39 kDa.

Positive Controls: rat skeletal muscle extract: sc-364810.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



ALP (H-157): sc-98652. Western blot analysis of ALP expression in rat skeletal muscle tissue extract.



ALP (H-157): sc-98652. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic and membrane staining of myocytes.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **ALP (B-10): sc-365765** or **ALP (C-8): sc-373737**, our highly recommended monoclonal alternatives to ALP (H-157).