

CRISP-2 (M-75): sc-98667

BACKGROUND

Cysteine-rich secretory proteins (CRISPs) represent a family of evolutionarily conserved proteins which may play a role in the innate immune system and are transcriptionally regulated by androgens in several tissues. CRISP-1 coats the postacrosomal region of sperm heads as they pass through the epididymis. CRISP-1 is found in all regions of the epididymis, ductus deferens, seminal plasma and sperm. CRISP-2, also known as testis-specific protein TPX1 or cancer/testis antigen 36 (CT36), is a 243 amino acid secreted protein. Expressed in the testis and epididymis, CRISP-2 is thought to be involved in calcium fluxes during sperm capacitation by regulating certain ion channels' activity. CRISP-3 is expressed in pancreas and prostate tissues, and along with CRISP-1, is expressed in saliva. The gene that encodes CRISP-3 is an early response gene that may participate in the pathophysiology of the autoimmune lesions of Sjogren's syndrome.

REFERENCES

1. Busso, D., Cohen, D.J., Hayashi, M., Kasahara, M. and Cuasnicú, P.S. 2005. Human testicular protein TPX1/CRISP-2: localization in spermatozoa, fate after capacitation and relevance for gamete interaction. *Mol. Hum. Reprod.* 11: 299-305.
2. Du, Y., Huang, X., Li, J., Hu, Y., Zhou, Z. and Sha, J. 2006. Human testis-specific protein 1 expression in human spermatogenesis and involvement in the pathogenesis of male infertility. *Fertil. Steril.* 85: 1852-1854.
3. Gibbs, G.M., Scanlon, M.J., Swarbrick, J., Curtis, S., Gallant, E., Dulhunty, A.F. and O'Bryan, M.K. 2006. The cysteine-rich secretory protein domain of TPX1 is related to ion channel toxins and regulates ryanodine receptor Ca^{2+} signaling. *J. Biol. Chem.* 281: 4156-4163.
4. Hamann, H., Jude, R., Sieme, H., Mertens, U., Töpfer-Petersen, E., Distl, O. and Leeb, T. 2007. A polymorphism within the equine CRISP-3 gene is associated with stallion fertility in Hanoverian warmblood horses. *Anim. Genet.* 38: 259-264.
5. Busso, D., Goldweic, N.M., Hayashi, M., Kasahara, M. and Cuasnicú, P.S. 2007. Evidence for the involvement of testicular protein CRISP-2 in mouse sperm-egg fusion. *Biol. Reprod.* 76: 701-708.

CHROMOSOMAL LOCATION

Genetic locus: Crisp2 (mouse) mapping to 17 B2.

SOURCE

CRISP-2 (M-75) is a rabbit polyclonal antibody raised against amino acids 24-98 mapping near the N-terminus of CRISP-2 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CRISP-2 (M-75) is recommended for detection of CRISP-2 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CRISP-2 siRNA (m): sc-77025, CRISP-2 shRNA Plasmid (m): sc-77025-SH and CRISP-2 shRNA (m) Lentiviral Particles: sc-77025-V.


Molecular Weight of CRISP-2: 27 kDa.

Positive Controls: mouse epididymus tissue extract or rat epididymus tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CRISP-2 (M-75): sc-98667. Western blot analysis of CRISP-2 expression in mouse epididymus (A) and rat epididymus (B) tissue extracts.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **CRISP-2 (D-10): sc-390914** or **CRISP-2 (E-12): sc-377391**, our highly recommended monoclonal alternatives to CRISP-2 (M-75).