

JMJD2C (H-300): sc-98678

BACKGROUND

JMJD2C (Jumonji domain containing 2C), also known as GASC1, KDM4C or JHDM3C, is a nuclear protein that belongs to the Jumonji domain 2 (JMJD2) family of histone demethylases. Functioning as a trimethylation-specific demethylase, JMJD2C demethylates specific lysine residues of Histone H3, thereby converting the trimethylated Histone H3 to its dimethylated form and playing a central role in the histone code. Through its ability to modify histones, JMJD2C increases the rate of cell proliferation and promotes the expression of a variety of proteins. JMJD2C binds iron as a cofactor and contains two Tudor domains through which it interacts with methylated histones. Overexpression of JMJD2C is associated with esophageal squamous cell carcinoma, suggesting a possible role for JMJD2C in carcinogenesis. Two isoforms of JMJD2C exist due to alternative splicing events.

REFERENCES

1. Yang, Z.Q., et al. 2000. Identification of a novel gene, GASC1, within an amplicon at 9p23-24 frequently detected in esophageal cancer cell lines. *Cancer Res.* 60: 4735-4739.
2. Katoh, M. and Katoh, M. 2004. Identification and characterization of JMJD2 family genes *in silico*. *Int. J. Oncol.* 24: 1623-1628.
3. Cloos, P.A., et al. 2006. The putative oncogene GASC1 demethylates tri- and dimethylated Lysine 9 on Histone H3. *Nature* 442: 307-311.
4. Whetstine, J.R., et al. 2006. Reversal of histone lysine trimethylation by the JMJD2 family of histone demethylases. *Cell* 125: 467-481.
5. Loh, Y.H., et al. 2007. JMJD1A and JMJD2C Histone H3 Lys 9 demethylases regulate self-renewal in embryonic stem cells. *Genes Dev.* 21: 2545-2557.
6. Katoh, Y. and Katoh, M. 2007. Comparative integromics on JMJD2A, JMJD2B and JMJD2C: preferential expression of JMJD2C in undifferentiated ES cells. *Int. J. Mol. Med.* 20: 269-273.
7. Wissmann, M., et al. 2007. Cooperative demethylation by JMJD2C and LSD1 promotes androgen receptor-dependent gene expression. *Nat. Cell Biol.* 9: 347-353.
8. Helias, C., et al. 2008. Polycythemia vera transforming to acute myeloid leukemia and complex abnormalities including 9p homogeneously staining region with amplification of MLLT3, JMJD2C, JAK2, and SMARCA2. *Cancer Genet. Cytogenet.* 180: 51-55.

CHROMOSOMAL LOCATION

Genetic locus: KDM4C (human) mapping to 9p24.1; Kdm4c (mouse) mapping to 4 C3.

SOURCE

JMJD2C (H-300) is a rabbit polyclonal antibody raised against amino acids 349-648 mapping within an internal region of JMJD2C of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

JMJD2C (H-300) is recommended for detection of JMJD2C of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for JMJD2C siRNA (h): sc-92765, JMJD2C siRNA (m): sc-146324, JMJD2C shRNA Plasmid (h): sc-92765-SH, JMJD2C shRNA Plasmid (m): sc-146324-SH, JMJD2C shRNA (h) Lentiviral Particles: sc-92765-V and JMJD2C shRNA (m) Lentiviral Particles: sc-146324-V.

Molecular Weight of JMJD2C: 120 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



MONOS
Satisfaction
Guaranteed

Try **JMJD2C (D-4): sc-515767**, our highly recommended monoclonal alternative to JMJD2C (H-300).