

FARP1 (H-300): sc-98722

BACKGROUND

FARP1 (FERM, RhoGEF (ARHGEF) and pleckstrin domain protein 1), also known as PLEKHC2 or CDEP, is a 1,045 amino acid protein that contains one FERM domain, one DH domain and 2 PH domains. Existing as multiple alternatively spliced isoforms that are expressed in fetal heart, brain and spleen, as well as in adult lung, kidney and testis, FARP1 is thought to function as a Rho-guanine nucleotide exchange factor that may play a role in linking the cell membrane to the cytoskeleton. The gene encoding FARP1 maps to human chromosome 13, which houses over 400 genes, such as BRCA2 and RB1, and comprises nearly 4% of the human genome. Trisomy 13, also known as Patau syndrome, is deadly and the few who survive past one year suffer from permanent neurologic defects, difficulty eating and vulnerability to serious respiratory infections.

CHROMOSOMAL LOCATION

Genetic locus: FARP1 (human) mapping to 13q32.2; Farp1 (mouse) mapping to 14 E5.

SOURCE

FARP1 (H-300) is a rabbit polyclonal antibody raised against amino acids 746-1045 mapping at the C-terminus of FARP1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

FARP1 (H-300) is recommended for detection of FARP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

FARP1 (H-300) is also recommended for detection of FARP1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for FARP1 siRNA (h): sc-77315, FARP1 siRNA (m): sc-77316, FARP1 shRNA Plasmid (h): sc-77315-SH, FARP1 shRNA Plasmid (m): sc-77316-SH, FARP1 shRNA (h) Lentiviral Particles: sc-77315-V and FARP1 shRNA (m) Lentiviral Particles: sc-77316-V.


Molecular Weight of FARP1: 119/122 kDa.

Positive Controls: F9 cell lysate: sc-2245, HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



FARP1 (H-300): sc-98722. Western blot analysis of FARP1 expression in c4 (A) and Jurkat (B) whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **FARP1 (2D4): sc-293249**, our highly recommended monoclonal alternative to FARP1 (H-300).