

Wnt-2b (H-39): sc-98737

BACKGROUND

The Wnt family of protooncogenes consists of at least 13 known members which encode secreted signaling proteins that are involved in oncogenesis and several other developmental processes, such as regulation of cell fate and embryogenesis. Wnt-2b (wingless-type MMTV integration site family, member 2B), also known as WNT13 or XWNT2, is a 391 amino acid protein that is secreted into the extracellular matrix and belongs to the Wnt family. Expressed as two isoforms, the first of which is present primarily in adult tissues and the second of which is present primarily in fetal tissues, Wnt-2b functions as a ligand for transmembrane receptor proteins and is thought to play a role in tissue differentiation and development. Wnt-2b is expressed in a variety of cancer cell lines, suggesting a role in tumorigenesis.

REFERENCES

- Wolda, S.L. and Moon, R.T. 1992. Cloning and developmental expression in *Xenopus laevis* of seven additional members of the Wnt family. *Oncogene* 7: 1941-1947.
- Smolich, B.D., et al. 1993. Wnt family proteins are secreted and associated with the cell surface. *Mol. Biol. Cell* 4: 1267-1275.
- Katoh, M., et al. 1996. Cloning, expression and chromosomal localization of Wnt-13, a novel member of the Wnt gene family. *Oncogene* 13: 873-876.
- Katoh, M., et al. 2000. Alternative splicing of the Wnt-2b/Wnt-13 gene. *Biochem. Biophys. Res. Commun.* 275: 209-216.
- Ricken, A., et al. 2002. Wnt signaling in the ovary: identification and compartmentalized expression of Wnt-2, Wnt-2b, and frizzled-4 mRNAs. *Endocrinology* 143: 2741-2749.
- Katoh, M. 2005. Wnt-2b: comparative integromics and clinical applications. *Int. J. Mol. Med.* 16: 1103-1108.
- Katoh, M. 2005. Molecular evolution of Wnt-2b orthologs. *Int. J. Oncol.* 26: 1135-1139.
- Struewing, I.T., et al. 2006. Mitochondrial and nuclear forms of Wnt-13 are generated via alternative promoters, alternative RNA splicing and alternative translation start sites. *J. Biol. Chem.* 281: 7282-7293.
- Ober, E.A., et al. 2006. Mesodermal Wnt-2b signalling positively regulates liver specification. *Nature* 442: 688-691.

CHROMOSOMAL LOCATION

Genetic locus: WNT2B (human) mapping to 1p13.2; Wnt2b (mouse) mapping to 3 F2.2.

SOURCE

Wnt-2b (H-39) is a rabbit polyclonal antibody raised against amino acids 263-301 mapping within an internal region of Wnt-2b of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Wnt-2b (H-39) is recommended for detection of Wnt-2b of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Wnt-2b (H-39) is also recommended for detection of Wnt-2b in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for Wnt-2b siRNA (h): sc-78902, Wnt-2b siRNA (m): sc-155356, Wnt-2b shRNA Plasmid (h): sc-78902-SH, Wnt-2b shRNA Plasmid (m): sc-155356-SH, Wnt-2b shRNA (h) Lentiviral Particles: sc-78902-V and Wnt-2b shRNA (m) Lentiviral Particles: sc-155356-V.

Molecular Weight (predicted) of Wnt-2b: 44 kDa.

Molecular Weight (observed) of Wnt-2b: 38 kDa.

Positive Controls: mouse brain extract: sc-2253 or Jurkat whole cell lysate: sc-2204.

DATA



Wnt-2b (H-39): sc-98737. Western blot analysis of Wnt-2b expression in mouse brain tissue extract.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.