



DC-STAMP (M-91): sc-98769

BACKGROUND

Bone homeostasis is accomplished by the balance of osteoblast and osteoclast activity. DC-STAMP (DC-specific transmembrane protein), also known as Transmembrane 7 superfamily member 4, is a 470 amino acid multi-pass membrane protein that regulates osteoclast and macrophage cell fusion. With localization at the cell surface, DC-STAMP is highly expressed in dendritic cells of the lung, kidney, liver and lymph nodes and is expressed at lower levels in spleen, bone marrow, leukocytes and pancreas. Mice that lack the gene encoding DC-STAMP exhibit upregulation of bone formation by osteoblasts and decreased bone reabsorption, ultimately leading to an increase in bone mass. Alternatively, mice that overexpress the DC-STAMP gene show decreased osteoblast activity and bone mass, due to accelerated cell-cell fusion of osteoclasts. This evidence suggests that DC-STAMP plays an essential role in osteoclastogenesis.

REFERENCES

- Hartgers, F.C., et al. 2000. DC-STAMP, a novel multimembrane-spanning molecule preferentially expressed by dendritic cells. *Eur. J. Immunol.* 30: 3585-3590.
- Staege, H., et al. 2001. Two novel genes FIND and LIND differentially expressed in deactivated and Listeria-infected human macrophages. *Immunogenetics* 53: 105-113.
- Hartgers, F.C., et al. 2001. Genomic organization, chromosomal localization, and 5' upstream region of the human DC-STAMP gene. *Immunogenetics* 53: 145-149.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605933. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Yagi, M., et al. 2005. DC-STAMP is essential for cell-cell fusion in osteoclasts and foreign body giant cells. *J. Exp. Med.* 202: 345-351.
- Iwasaki, R., et al. 2008. Cell fusion in osteoclasts plays a critical role in controlling bone mass and osteoblastic activity. *Biochem. Biophys. Res. Commun.* 377: 899-904.
- Sawatani, Y., et al. 2008. The role of DC-STAMP in maintenance of immune tolerance through regulation of dendritic cell function. *Int. Immunol.* 20: 1259-1268.
- Eleveld-Trancikova, D., et al. 2008. The DC-derived protein DC-STAMP influences differentiation of myeloid cells. *Leukemia* 22: 455-459.
- Jansen, B.J., et al. 2009. OS9 interacts with DC-STAMP and modulates its intracellular localization in response to TLR ligation. *Mol. Immunol.* 46: 505-515.

CHROMOSOMAL LOCATION

Genetic locus: TM7SF4 (human) mapping to 8q22.3; Tm7sf4 (mouse) mapping to 15 B3.1.

SOURCE

DC-STAMP (M-91) is a rabbit polyclonal antibody raised against amino acids 119-209 mapping within an internal region of DC-STAMP of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

DC-STAMP (M-91) is recommended for detection of DC-STAMP of mouse, rat and, to a lesser extent, human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DC-STAMP siRNA (h): sc-77688, DC-STAMP siRNA (m): sc-142887, DC-STAMP shRNA Plasmid (h): sc-77688-SH, DC-STAMP shRNA Plasmid (m): sc-142887-SH, DC-STAMP shRNA (h) Lentiviral Particles: sc-77688-V and DC-STAMP shRNA (m) Lentiviral Particles: sc-142887-V.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.