

HIF PHD2 (H-49): sc-98789

BACKGROUND

Prolyl hydroxylase domain proteins HIF PHD1, HIF PHD2 and HIF PHD3 (also designated EGLN2, EGLN1 and EGLN3, respectively) can hydroxylate HIF- α subunits. Hypoxia-inducible factor (HIF) is a transcriptional regulator important in several aspects of oxygen homeostasis. The prolyl hydroxylases catalyze the posttranslational formation of 4-hydroxyproline in HIF- α proteins. HIF PHD1, which is widely expressed, with highest levels of expression in testis, functions as a cellular oxygen sensor and is important in cell growth regulation. HIF PHD1 can localize to the nucleus or the cytoplasm and is also detected in hormone responsive tissues, such as normal and cancerous mammary, ovarian and prostate epithelium. HIF PHD1 is encoded by EGLN2, which maps to chromosome 19q13.3. HIF PHD2 is regarded as the main cellular oxygen sensor, as RNA interference against HIF PHD2, but not HIF PHD1 or HIF PHD3, is enough to stabilize HIF-1 α in normoxia. HIF PHD2, a direct HIF target gene, is expressed mainly in skeletal muscle, heart, kidney and brain. HIF PHD3 may play a role in the regulation of cell growth in muscle cells and in apoptosis in neuronal tissue. HIF PHD3 is widely expressed, although the highest levels can be detected in placenta and heart.

CHROMOSOMAL LOCATION

Genetic locus: EGLN1 (human) mapping to 1q42.2; EglN1 (mouse) mapping to 8 E2.

SOURCE

HIF PHD2 (H-49) is a rabbit polyclonal antibody raised against amino acids 240-288 mapping within an internal region of HIF PHD2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

HIF PHD2 (H-49) is recommended for detection of HIF PHD2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HIF PHD2 (H-49) is also recommended for detection of HIF PHD2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for HIF PHD2 siRNA (h): sc-45537, PHD2 siRNA (m): sc-45538, HIF PHD2 shRNA Plasmid (h): sc-45537-SH, PHD2 shRNA Plasmid (m): sc-45538-SH, HIF PHD2 shRNA (h) Lentiviral Particles: sc-45537-V and PHD2 shRNA (m) Lentiviral Particles: sc-45538-V.


Molecular Weight of HIF PHD2: 46 kDa.

Positive Controls: HIF PHD2 (m): 293T Lysate: sc-120777, Rat heart extract: sc-2393 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



HIF PHD2 (H-49): sc-98789. Western blot analysis of HIF PHD2 expression in HeLa (A), A549 (B), Hep G2 (C) and MCF7 (D) whole cell lysates and rat heart tissue extract (E).



HIF PHD2 (H-49): sc-98789. Western blot analysis of HIF PHD2 expression in non-transfected: sc-117752 (A) and mouse HIF PHD2 transfected: sc-120777 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **HIF PHD2 (H-8): sc-271835**, our highly recommended monoclonal alternative to HIF PHD2 (H-49).