

CD109 (H-91): sc-98793

BACKGROUND

CD109 is a glycosylphosphatidylinositol (GPI)-linked cell surface glycoprotein. It is a member of the α -macroglobulin/C3, C4, C5 family of thioester-containing proteins. The protein is expressed by CD34⁺ acute myeloid leukemia cell lines, activated T lymphoblasts, activated platelets, T cell lines, endothelial cells, lung and esophageal squamous cell carcinomas, testis and T cell lines. It has all the characteristics of a cancer-testis antigen. CD109 carries the platelet-specific Gov antigen system which is involved in platelet transfusion refraction, neonatal alloimmune thrombocytopenia and posttransfusion purpura.

REFERENCES

1. Sasaki, R., Takaku, F., Sakamoto, S. and Kanoh, Y. 1979. Terminal deoxynucleotidyl transferase activity and B cell markers in chronic myelogenous leukemia blast crisis. *Acta Haematol.* 62: 143-147.
2. Kelton, J.G., Smith, J.W., Horsewood, P., Humbert, J.R., Hayward, C.P. and Warkentin, T.E. 1990. Gova/b alloantigen system on human platelets. *Blood* 75: 2172-2176.
3. Lin, M., Sutherland, D.R., Horsfall, W., Totty, N., Yeo, E., Nayar, R., Wu, X.F. and Schuh, A.C. 2002. Cell surface antigen CD109 is a novel member of the α_2 -macroglobulin/C3, C4, C5 family of thioester-containing proteins. *Blood* 99: 1683-1691.
4. Schuh, A.C., Watkins, N.A., Nguyen, Q., Harmer, N.J., Lin, M., Prosper, J.Y., Campbell, K., Sutherland, D.R., Metcalfe, P., Horsfall, W., Ouwehand and W.H. 2002. A tyrosine703serine polymorphism of CD109 defines the Gov platelet alloantigens. *Blood* 99: 1692-1698.
5. Giesert, C., Marxer, A., Sutherland, D.R., Schuh, A.C., Kanz, L., Buhring and H.J. 2003. Antibody W7C5 defines a CD109 epitope expressed on CD34⁺ and CD34⁻ hematopoietic and mesenchymal stem cell subsets. *Ann. N.Y. Acad. Sci.* 996227-996230.
6. Solomon, K.R., Sharma, P., Chan, M., Morrison, P.T. and Finberg, R.W. 2004. CD109 represents a novel branch of the α_2 -macroglobulin/complement gene family. *Gene* 327: 171-183.

CHROMOSOMAL LOCATION

Genetic locus: CD109 (human) mapping to 6q13; Cd109 (mouse) mapping to 9 E1.

SOURCE

CD109 (H-91) is a rabbit polyclonal antibody raised against amino acids 957-1041 mapping within an internal region of CD109 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

CD109 (H-91) is recommended for detection of CD109 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CD109 (H-91) is also recommended for detection of CD109 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for CD109 siRNA (h): sc-44950, CD109 siRNA (m): sc-44951, CD109 shRNA Plasmid (h): sc-44950-SH, CD109 shRNA Plasmid (m): sc-44951-SH, CD109 shRNA (h) Lentiviral Particles: sc-44950-V and CD109 shRNA (m) Lentiviral Particles: sc-44951-V.


Molecular Weight of CD109: 170 kDa.

Positive Controls: rat testis extract: sc-2400, mouse testis extract: sc-2405 or human platelet whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



CD109 (H-91): sc-98793. Western blot analysis of CD109 expression in human platelet extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **CD109 (C-9): sc-271085** or **CD109 (H-7): sc-365780**, our highly recommended monoclonal alternatives to CD109 (H-91).