

UHRF1 (M-132): sc-98817

BACKGROUND

UHRF1 (ubiquitin-like, containing PHD and RING finger domains, 1), also known as Np95 (nuclear zinc finger protein 95), ICBP90 (inverted CCAAT box-binding protein of 90 kDa) or RNF106, is a transcription and cell cycle regulator belonging to the RING-finger type E3 ubiquitin ligase subfamily. UHRF1 is expressed in bone marrow, thymus, heart, testis and lung, and contains one PHD-type zinc finger, a ubiquitin-like domain, two RING-type zinc fingers and one YDG/SRA domain. Localizing to the nucleus, UHRF1 is believed to function as an E3 ubiquitin-protein ligase that accepts a ubiquitin residue from an E2 ubiquitin-conjugating enzyme and immediately transfers that residue to a protein that is targeted for degradation. By mediating ubiquitination, UHRF1 plays an important role in cellular proliferation. In addition, UHRF1 directly interacts with Dnmt1 (a maintenance DNA methyltransferase) and is required for the stable association of Dnmt1 with chromatin. UHRF1 is overexpressed in cancer cells, suggesting a possible role in carcinogenesis.

REFERENCES

- Hopfner, R., et al. 2000. ICBP90, a novel human CCAAT binding protein, involved in the regulation of topoisomerase II α expression. *Cancer Res.* 60: 121-128.
- Muto, M., et al. 2002. Targeted disruption of Np95 gene renders murine embryonic stem cells hypersensitive to DNA damaging agents and DNA replication blocks. *J. Biol. Chem.* 277: 34549-34555.
- Bonapace, I.M., et al. 2002. Np95 is regulated by E1A during mitotic reactivation of terminally differentiated cells and is essential for S phase entry. *J. Cell Biol.* 157: 909-914.
- Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607990. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Uhrf1 (mouse) mapping to 17 D.

SOURCE

UHRF1 (M-132) is a rabbit polyclonal antibody raised against amino acids 586-717 mapping near the C-terminus of UHRF1 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

UHRF1 (M-132) is recommended for detection of UHRF1 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for UHRF1 siRNA (m): sc-155976, UHRF1 shRNA Plasmid (m): sc-155976-SH and UHRF1 shRNA (m) Lentiviral Particles: sc-155976-V.


Molecular Weight of UHRF1: 90 kDa.

Positive Controls: mouse thymus extract: sc-2406.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



UHRF1 (M-132): sc-98817. Western blot analysis of UHRF1 expression in mouse thymus tissue extract.

SELECT PRODUCT CITATIONS

- Pacaud, R., et al. 2014. The DNMT1/PCNA/UHRF1 disruption induces tumorigenesis characterized by similar genetic and epigenetic signatures. *Sci. Rep.* 4: 4230.



Try **UHRF1 (H-8): sc-373750** or **UHRF1 (G-2): sc-166898**, our highly recommended monoclonal alternatives to UHRF1 (M-132). Also, for AC, HRP, FITC, PE, Alexa Fluor[®] 488 and Alexa Fluor[®] 647 conjugates, see **UHRF1 (H-8): sc-373750**.