

PRAM-1 (M-300): sc-98880

BACKGROUND

Complete remission of acute promyelocytic leukemia can be achieved by treating patients with retinoic acid, and PML-RAR α (promyelocytic leukemia-retinoic acid receptor α fusion protein) plays a major role in mediating retinoic acid effects in leukemia cells. The retinoic acid-induced gene, PRAM-1 (PML-RAR α target gene encoding an adaptor molecule 1) encodes an adaptor protein which is expressed and modulated during normal human myelopoiesis. PRAM-1 expression is hindered by expression of PML-RAR α . The 718 amino acid PRAM-1 protein contains 8 N-terminal proline-rich repeats and several proline residues that are clustered as type I or type II SH3 recognition motifs. PRAM-1 demonstrates expression in hematopoietic tissues and lung.

REFERENCES

- Moog-Lutz, C., et al. 2001. PRAM-1 is a novel adaptor protein regulated by retinoic acid (RA) and promyelocytic leukemia (PML)-RA receptor α in acute promyelocytic leukemia cells. *J. Biol. Chem.* 276: 22375-22381.
- Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 606466. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Clemens, R.A. et al. 2004. PRAM-1 is required for optimal integrin-dependent neutrophil function. *Mol. Cell. Biol.* 24: 10923-10932.
- Denis, F.M. et al. 2005. PRAM-1 potentiates arsenic trioxide-induced JNK activation. *J. Biol. Chem.* 280: 9043-9048.
- Heuer, K. et al. 2006. Lipid-binding HSH3 domains in immune cell adapter proteins. *J. Mol. Biol.* 361: 94-104.
- Susic, D. et al. 2006. Cardiovascular effects of nonproteolytic activation of prorenin. *Hypertension* 48: E113.
- Ghaffari, S.H. et al. 2006. Real-time PCR analysis of PML-RAR α in newly diagnosed acute promyelocytic leukaemia patients treated with arsenic trioxide as a front-line therapy. *Ann. Oncol.* 17: 1553-1559.

CHROMOSOMAL LOCATION

Genetic locus: PRAM1 (human) mapping to 19p13.2; Pram1 (mouse) mapping to 17 B1.

SOURCE

PRAM-1 (M-300) is a rabbit polyclonal antibody raised against amino acids 284-580 mapping within an internal region of PRAM-1 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

PRAM-1 (M-300) is recommended for detection of PRAM-1 of mouse, rat and, to a lesser extent, human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PRAM-1 siRNA (h): sc-61393, PRAM-1 siRNA (m): sc-61394, PRAM-1 shRNA Plasmid (h): sc-61393-SH, PRAM-1 shRNA Plasmid (m): sc-61394-SH, PRAM-1 shRNA (h) Lentiviral Particles: sc-61393-V and PRAM-1 shRNA (m) Lentiviral Particles: sc-61394-V.


Molecular Weight of PRAM-1: 97 kDa.

Positive Controls: PRAM-1 (h3): 293T Lysate: sc-159247.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



PRAM-1 (M-300): sc-98880. Western blot analysis of PRAM-1 expression in non-transfected: sc-117752 (A) and human PRAM-1 transfected: sc-159247 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **PRAM-1 (D-11): sc-166267** or **PRAM-1 (A-5): sc-376451**, our highly recommended monoclonal alternatives to PRAM-1 (M-300).