

tetranectin (H-100): sc-98886

BACKGROUND

tetranectin is a homotrimeric glycoprotein present in plasma and various tissue locations that binds to calcium, heparin and plasminogen kringle 4. Tetranectin may play a prominent role in tissue remodeling as well as in the regulation of proteolytic processes via its binding and indirect activation of plasminogen. tetranectin is found in the extracellular matrix (ECM) of certain carcinomas, but is not present in the ECM of normal tissues. Extracellular proteolysis is an important factor in the ability of malignant cells to penetrate normal tissues and metastasize. Decreased plasma tetranectin or increased tetranectin in stroma of cancers correlates with cancer progression and a grim prognosis. tetranectin may also influence cancer growth by altering activities of plasminogen or the plasminogen fragment, Angiostatin, which inhibits tumor cell proliferation.

REFERENCES

- Holtet, T.L., Graversen, J.H., Clemmensen, I., Thøgersen, H.C. and Etzerodt, M. 1997. Tetranectin, a trimeric plasminogen-binding C-type lectin. *Protein Sci.* 6: 1511-1515.
- Thougaard, A.V., Høgdall, C.K., Kjaer, S.K., Blaakaer, J., Jaliashvili, I. and Christiansen, M. 1998. Determination of serum tetranectin: technical and clinical evaluation of three sandwich immunoassays. *Clin. Chim. Acta* 276: 19-34.
- Høgdall, C.K. 1998. Human tetranectin: methodological and clinical studies. *APMIS Suppl.* 86: 1-31.
- Kamper, E.F., Papaphilis, A.D., Angelopoulou, M.K., Kopeikina, L.T., Siakantaris, M.P., Pangalis, G.A. and Stavridis, J.C. 1999. Serum levels of tetranectin, intercellular adhesion molecule-1 and interleukin-10 in B-chronic lymphocytic leukemia. *Clin. Biochem.* 32: 639-645.
- Lorentsen, R.H., Graversen, J.H., Caterer, N.R., Thøgersen, H.C. and Etzerodt, M. 2000. The heparin-binding site in tetranectin is located in the N-terminal region and binding does not involve the carbohydrate recognition domain. *Biochem. J.* 347: 83-87.
- Caterer, N.R., Graversen, J.H., Jacobsen, C., Moestrup, S.K., Sigurskjold, B.W., Etzerodt, M. and Thøgersen, H.C. 2002. Specificity determinants in the interaction of apolipoprotein A kringles with tetranectin and LDL. *Biol. Chem.* 383: 1743-1750.
- Nielbo, S., Thomsen, J.K., Graversen, J.H., Jensen, P.H., Etzerodt, M., Poulsen, F.M. and Thøgersen, H.C. 2004. Structure of the plasminogen kringle 4 binding calcium-free form of the C-type lectin-like domain of tetranectin. *Biochemistry* 43: 8636-8643.

CHROMOSOMAL LOCATION

Genetic locus: CLEC3B (human) mapping to 3p21.31; Clec3b (mouse) mapping to 9 F4.

SOURCE

tetranectin (H-100) is a rabbit polyclonal antibody raised against amino acids 103-202 mapping at the C-terminus of tetranectin of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

tetranectin (H-100) is recommended for detection of tetranectin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

tetranectin (H-100) is also recommended for detection of tetranectin in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for tetranectin siRNA (h): sc-61675, tetranectin siRNA (m): sc-61676, tetranectin shRNA Plasmid (h): sc-61675-SH, tetranectin shRNA Plasmid (m): sc-61676-SH, tetranectin shRNA (h) Lentiviral Particles: sc-61675-V and tetranectin shRNA (m) Lentiviral Particles: sc-61676-V.

Molecular Weight of tetranectin: 20 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfation
Guaranteed

Try **tetranectin (F-8): sc-376940** or **tetranectin (10E3): sc-80594**, our highly recommended monoclonal alternatives to tetranectin (H-100).