

NPY5-R (H-40): sc-98935

BACKGROUND

Pancreatic polypeptide (PP), neuropeptide Y (NPY), and peptide YY (PYY) are related 36-amino acid hormones. A number of structurally related receptors for these peptides have been isolated, NPY1-R, NPY2-R, NPY3-R, NPY4-R, NPY5-R and NPY6-R. NPY4-R is expressed in several human tissues, including brain, coronary artery and ileum. NPY4-R maps to human chromosome 10q11.2-q21.2. NPY5-R, isolated from rat hypothalamus, encodes a 456 amino acid protein with less than 35% overall identity to known Y-type receptors. The human NPY5-R sequence is nearly identical to, but in the opposite orientation from, that of the human NPY1-R sequence. NPY5-R localizes to the paraventricular hypothalamic nucleus, the lateral hypothalamus and other locations consistent with a role in the control of feeding behavior. The gene which encodes NPY5-R maps to human chromosome 4q32.2. NPY6-R is abundantly expressed in human heart and skeletal muscle and the gene which encodes NPY6-R maps to human chromosome 5q31.

REFERENCES

1. Bard, J.A., et al. 1995. Cloning and functional expression of a human Y4 subtype receptor for pancreatic polypeptide, neuropeptide Y, and peptide YY. *J. Biol. Chem.* 270: 26762-26765.
2. Gerald, C., et al. 1996. A receptor subtype involved in neuropeptide-Y-induced food intake. *Nature* 382: 168-171.
3. Hu, Y., et al. 1996. Identification of a novel hypothalamic neuropeptide Y receptor associated with feeding behavior. *J. Biol. Chem.* 271: 26315-26319.
4. Matsumoto, M., et al. 1996. Inactivation of a novel neuropeptide Y/peptide YY receptor gene in primate species. *J. Biol. Chem.* 271: 27217-27220.
5. Herzog, H., et al. 1997. Overlapping gene structure of the human neuropeptide Y receptor subtypes Y1 and Y5 suggests coordinate transcriptional regulation. *Genomics* 41: 315-319.
6. Lutz, C.M., et al. 1997. Neuropeptide Y receptor genes mapped in human and mouse: receptors with high affinity for pancreatic polypeptide are not clustered with receptors specific for neuropeptide Y and peptide YY. *Genomics* 46: 287-290.
7. Darby, K., et al. 1997. Assignment of the Y-4 receptor gene (PPYR1) to human chromosome 10q11.2 and mouse chromosome 14. *Genomics* 46: 513-515.

CHROMOSOMAL LOCATION

Genetic locus: NPY5R (human) mapping to 4q32.2; Npy5r (mouse) mapping to 8 B3.2.

SOURCE

NPY5-R (H-40) is a rabbit polyclonal antibody raised against amino acids 391-430 mapping near the C-terminus of NPY5-R of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

NPY5-R (H-40) is recommended for detection of NPY5-R of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

NPY5-R (H-40) is also recommended for detection of NPY5-R in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for NPY5-R siRNA (h): sc-42105, NPY5-R siRNA (m): sc-42106, NPY5-R shRNA Plasmid (h): sc-42105-SH, NPY5-R shRNA Plasmid (m): sc-42106-SH, NPY5-R shRNA (h) Lentiviral Particles: sc-42105-V and NPY5-R shRNA (m) Lentiviral Particles: sc-42106-V.

Molecular Weight of NPY5-R: 57 kDa.

Positive Controls: MIA PaCa-2 cell lysate: sc-2285, SK-N-SH cell lysate: sc-2410 or Ramos cell lysate: sc-2216.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **NPY5-R (H-4): sc-137167**, our highly recommended monoclonal alternative to NPY5-R (H-40).