

## PKA $\gamma$ cat (H-55): sc-98950

### BACKGROUND

The second messenger cyclic AMP (cAMP) mediates diverse cellular responses to external signals such as proliferation, ion transport, regulation of metabolism and gene transcription by activation of the cAMP-dependent protein kinase (cAPK or PKA). Activation of PKA occurs when cAMP binds to the two regulatory subunits of the tetrameric PKA holoenzyme resulting in release of active catalytic subunits. Three catalytic (C) subunits have been identified, designated C $\alpha$ , C $\beta$  and C $\gamma$ , that each represent specific gene products. C $\alpha$  and C $\beta$  are closely related (93% amino acid sequence similarity), whereas C $\gamma$  displays 83% and 79% similarity to C $\alpha$  and C $\beta$ , respectively. Activation of transcription upon elevation of cAMP levels results from translocation of PKA to the nucleus where it phosphorylates the transcription factor cAMP response element binding protein (CREB) on serine 133 which in turn leads to TFIIB binding to TATA-box-binding protein TBP1, thus linking phospho-CREB to the Pol II transcription initiation complex.

### REFERENCES

1. Beavo, J.A., Bechtel, P.J. and Krebs, E.G. 1974. Activation of protein kinase by physiological concentrations of cyclic AMP. *Proc. Natl. Acad. Sci. USA* 71: 3580-3583.
2. Krebs, E.G. and Beavo, J.A. 1980. Phosphorylation and dephosphorylation of enzymes. *Ann. Rev. Biochem.* 48: 923-959.
3. Maldonado, F. and Hanks, SK. 1988. cAMP-dependent protein kinase,  $\alpha$ -catalytic subunit. *Nucl. Acids Res.* 16: 8189-8190.
4. Gonzalez, G.A. and Montminy, M.R. 1989. Cyclic AMP stimulates somatostatin gene transcription by phosphorylation of CREB at serine 133. *Cell* 59: 675-680.
5. Beebe, S.J., Oyen, O., Sandberg, M., Frøysa, A., Hansson, V. and Jahnsen, T. 1990. cAMP-dependent protein kinase,  $\beta$ -catalytic subunit. *Mol. Endocrinol.* 4: 465-475.
6. Meinkoth, J.L., Alberts, A.S., Went, W., Fantozzi, D., Taylor, S.S., Hagiwara, M., Montminy, M. and Feramisco, J.R. 1993. Signal transduction through the cAMP-dependent protein kinase. *Mol. Cell. Biochem.* 127-128: 179-186.
7. Nordheim, A. 1994. CREB takes CBP to tango. *Nature* 370: 177-178.

### CHROMOSOMAL LOCATION

Genetic locus: PRKACG (human) mapping to 9q21.11.

### SOURCE

PKA $\gamma$  cat (H-55) is a rabbit polyclonal antibody raised against amino acids 6-55 mapping at the N-terminus of PKA $\gamma$  catalytic subunit of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### APPLICATIONS

PKA $\gamma$  cat (H-55) is recommended for detection of PKA $\gamma$  catalytic subunit of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with PKA $\alpha$  subunit.

Suitable for use as control antibody for PKA $\gamma$  cat siRNA (h): sc-36236, PKA $\gamma$  cat shRNA Plasmid (h): sc-36236-SH and PKA $\gamma$  cat shRNA (h) Lentiviral Particles: sc-36236-V.

Molecular Weight of PKA $\gamma$  cat: 40 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
 Satisfaction  
 Guaranteed

Try **PKA $\gamma$  cat (A-4): sc-514087**, our highly recommended monoclonal alternative to PKA $\gamma$  cat (H-55).