

RGS13 (H-135): sc-99132

BACKGROUND

The regulators of G protein signaling (RGS) proteins inhibit heterotrimeric G protein signaling. RGS proteins work by functioning as GTPase-activating proteins (which increase the GTPase activity of G protein α subunits) thereby driving G proteins into their inactive GDP-bound form. RGS13 (regulator of G protein signaling 13) localizes in membrane and nuclear fractions, and is expressed predominantly in tonsil, thymus, lymph node, lung and spleen tissues. Within the lymphoid compartment, highest levels of RGS13 have been found in resting CD19-positive (B cells). Unlike most RGS proteins, RGS13 has no recognizable domain other than the RGS box, but because of its prevalence in the immune system and lung and its ability to inhibit $G_{\alpha q}$, $G_{\alpha i}$ and cAMP generation, the function of RGS13 might be to modulate specific G protein-dependent signal transduction pathways in these regions.

REFERENCES

1. Johnson, E.N. and Druey, K.M. 2002. Functional characterization of the G protein regulator RGS13. *J. Biol. Chem.* 277: 16768-16774.
2. Shi, G.X., et al. 2002. RGS13 regulates germinal center B lymphocytes responsiveness to CXCL chemokine ligand (CXCL12 and CXCL13). *J. Immunol.* 169: 2507-2515.
3. Sierra, D.A., et al. 2002. Evolution of the regulators of G protein signaling multigene family in mouse and human. *Genomics* 79: 177-185.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607190. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Estes, J.D., et al. 2004. Follicular dendritic cell regulation of CXCR4-mediated germinal center CD4 T cell migration. *J. Immunol.* 173: 6169-6178.
6. Moratz, C., et al. 2004. Regulation of chemokine-induced lymphocyte migration by RGS proteins. *Meth. Enzymol.* 389: 15-32.
7. Han, J.I., et al. 2006. RGS1 and RGS13 mRNA silencing in a human B lymphoma line enhances responsiveness to chemoattractants and impairs desensitization. *J. Leukoc. Biol.* 79: 1357-1368.

CHROMOSOMAL LOCATION

Genetic locus: RGS13 (human) mapping to 1q31.2; Rgs13 (mouse) mapping to 1 F.

SOURCE

RGS13 (H-135) is a rabbit polyclonal antibody raised against amino acids 1-135 mapping at the N-terminus of RGS13 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

RGS13 (H-135) is recommended for detection of RGS13 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RGS13 (H-135) is also recommended for detection of RGS13 in additional species, including equine.

Suitable for use as control antibody for RGS13 siRNA (h): sc-61464, RGS13 siRNA (m): sc-61465, RGS13 shRNA Plasmid (h): sc-61464-SH, RGS13 shRNA Plasmid (m): sc-61465-SH, RGS13 shRNA (h) Lentiviral Particles: sc-61464-V and RGS13 shRNA (m) Lentiviral Particles: sc-61465-V.


Molecular Weight of RGS13: 19 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



RGS13 (H-135): sc-99132. Western blot analysis of RGS13 expression in mouse brain tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **RGS13 (G-7): sc-514590**, our highly recommended monoclonal alternative to RGS13 (H-135).