

MUTYH (H-44): sc-99211

BACKGROUND

MUTYH (mutY homolog (*E. coli*)) is a DNA glycosylase mismatch repair enzyme that in conjunction with mutM (OGG1), cleaves adenine residues paired with either oxidized (8-hydroxyguanines) or non-modified guanines in order to correct A/G and A/C mismatches. Repair of most modified and mis-paired bases in the genome is initiated by DNA glycosylases, which bind and cleave N-glycosyl bonds to initiate base excision repair. MUTYH is crucial for the avoidance of mutations resulting from oxidative DNA damage. Multiple N-terminal splice variants of MUTYH exist in mammalian cells. Increasing levels of MUTYH in A549 cells exposed to oxygen and infrared radiation leads to improvements in cell survival. Biallelic MUTYH germ-line mutations predispose humans to colorectal adenomas and carcinomas. MUTYH is abundant in neurons where mitochondrial genomes exposed to reactive oxygen species (ROS) that damage DNA must maintain integrity over the entire mammalian life span.

REFERENCES

- Hayashi, H., et al. 2002. Replication-associated repair of adenine: 8-oxoguanine mispairs by MYH. *Curr. Biol.* 12: 335-339.
- Englander, E.W., et al. 2002. Rat MYH, a glycosylase for repair of oxidatively damaged DNA, has brain-specific isoforms that localize to neuronal mitochondria. *J. Neurochem.* 83: 1471-1480.
- Halford, S.E., et al. 2003. Germline mutations but not somatic changes at the MYH locus contribute to the pathogenesis of unselected colorectal cancers. *Am. J. Pathol.* 162: 1545-1548.
- Lee, H.M., et al. 2004. Developmental changes in expression and subcellular localization of the DNA repair glycosylase, MYH, in the rat brain. *J. Neurochem.* 88: 394-400.
- Tao, H., et al. 2004. A novel splice-site variant of the base excision repair gene MYH is associated with production of an aberrant mRNA transcript encoding a truncated MYH protein not localized in the nucleus. *Carcinogenesis* 25: 1859-1866.
- Kim, C.J., et al. 2004. Genetic alterations of the MYH gene in gastric cancer. *Oncogene* 23: 6820-6822.
- Ma, H., et al. 2004. N-terminus of the rat adenine glycosylase MYH affects excision rates and processing of MYH-generated abasic sites. *Nucleic Acids Res.* 32: 4332-4339.
- Sieber, O.M., et al. 2004. MYH deficiency enhances intestinal tumorigenesis in multiple intestinal neoplasia (ApcMin⁺) mice. *Cancer Res.* 64: 8876-8881.
- Cardoso, J., et al. 2006. Chromosomal instability in MYH- and APC-mutant adenomatous polyps. *Cancer Res.* 66: 2514-2519.

CHROMOSOMAL LOCATION

Genetic locus: MUTYH (human) mapping to 1p34.1; Mutyh (mouse) mapping to 4 D1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

MUTYH (H-44) is a rabbit polyclonal antibody raised against amino acids 182-225 mapping within an internal region of MUTYH of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

MUTYH (H-44) is recommended for detection of MUTYH of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MUTYH (H-44) is also recommended for detection of MUTYH in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for MUTYH siRNA (h): sc-37407, MUTYH siRNA (m): sc-45816, MUTYH shRNA Plasmid (h): sc-37407-SH, MUTYH shRNA Plasmid (m): sc-45816-SH, MUTYH shRNA (h) Lentiviral Particles: sc-37407-V and MUTYH shRNA (m) Lentiviral Particles: sc-45816-V.

Molecular Weight of MUTYH: 65 kDa.

Positive Controls: HeLa nuclear extract: sc-2120, HeLa whole cell lysate: sc-2200 or SW480 nuclear extract: sc-2155.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.



Try **MUTYH (C-6): sc-374571**, our highly recommended monoclonal alternative to MUTYH (H-44).