

## HYPB (D-19): sc-99451

### BACKGROUND

HYPB (huntingtin yeast partner B), also known as SETD2 (SET domain-containing protein 2) or HIF1, is a 2,564 amino acid nuclear protein that contains one WW domain, one SET domain, one post-SET domain and one AWS domain and belongs to the huntingtin interacting protein family. Expressed ubiquitously, HYPB functions as a histone methyltransferase that is specific for the lysine-36 residue of Histone H3 which, once methylated, plays a role in transcriptional activation and is associated with active chromatin. Due to its role in Histone H3 methylation, HYPB is thought to be involved in the modulation of chromatin structure and may also bind to DNA promoters and interact with Pol II, thereby promoting transcription. HYPB may be associated with the pathogenesis of the neurodegenerative disorder Huntington's disease (HD), which is characterized by a loss of striatal neurons, leading to brain deterioration and, ultimately, death. Three isoforms of HYPB exist as a result of alternative splicing events.

### REFERENCES

1. Faber, P.W., Barnes, G.T., Srinidhi, J., Chen, J., Gusella, J.F. and MacDonald, M.E. 1998. Huntingtin interacts with a family of WW domain proteins. *Hum. Mol. Genet.* 7: 1463-1474.
2. Passani, L.A., Bedford, M.T., Faber, P.W., McGinnis, K.M., Sharp, A.H., Gusella, J.F., Vonsattel, J.P. and MacDonald, M.E. 2000. Huntingtin's WW domain partners in Huntington's disease post-mortem brain fulfill genetic criteria for direct involvement in Huntington's disease pathogenesis. *Hum. Mol. Genet.* 9: 2175-2182.
3. Rega, S., Stiewe, T., Chang, D.I., Pollmeier, B., Esche, H., Bardenheuer, W., Marquitan, G. and Pützer, B.M. 2001. Identification of the full-length huntingtin-interacting protein p231HBP/HYPB as a DNA-binding factor. *Mol. Cell. Neurosci.* 18: 68-79.
4. Sun, X.J., Wei, J., Wu, X.Y., Hu, M., Wang, L., Wang, H.H., Zhang, Q.H., Chen, S.J., Huang, Q.H. and Chen, Z. 2005. Identification and characterization of a novel human Histone H3 lysine 36-specific methyltransferase. *J. Biol. Chem.* 280: 35261-35271.
5. Li, M., Phatnani, H.P., Guan, Z., Sage, H., Greenleaf, A.L. and Zhou, P. 2005. Solution structure of the Set2-Rpb1 interacting domain of human Set2 and its interaction with the hyperphosphorylated C-terminal domain of Rpb1. *Proc. Natl. Acad. Sci. USA* 102: 17636-17641.

### CHROMOSOMAL LOCATION

Genetic locus: SETD2 (human) mapping to 3p21.31; Setd2 (mouse) mapping to 9 F2.

### SOURCE

HYPB (D-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HYPB of human origin.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-99451 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-99451 X, 100 µg/0.1 ml.

### APPLICATIONS

HYPB (D-19) is recommended for detection of HYPB of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HYPB (D-19) is also recommended for detection of HYPB in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for HYPB siRNA (h): sc-78103, HYPB shRNA Plasmid (h): sc-78103-SH and HYPB shRNA (h) Lentiviral Particles: sc-78103-V.

HYPB (D-19) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of HYPB precursor splice form 1: 285 kDa.

Molecular Weight of HYPB splice form 2: 192 kDa.


Molecular Weight of HYPB splice form 3: 176 kDa.

Positive Controls: Jurkat nuclear extract: sc-2132 or U-937 nuclear extract: sc-2156.

### DATA



HYPB (D-19): sc-99451. Western blot analysis of HYPB expression in Jurkat (A) and U-937 (B) nuclear extracts.



HYPB (D-19): sc-99451. Immunofluorescence staining of formalin-fixed HeLa cells showing nuclear localization. Kindly provided by Yang Xiang, Ph.D., Division of Newborn Medicine, Boston Children's Hospital, Cell Biology Department, Harvard Medical School.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.