

Intersectin-2 (H-16): sc-9953

BACKGROUND

Intersectin, which is also designated Ese1 or ITSN1, is a component of the cellular endocytic machinery. Intersectin is composed of two N-terminal Ese15 homology (EH) domains, a central highly charged region and five C-terminal SH3 domains, which all largely contribute to the association of Intersectin with other components of the endocytic pathway. The EH domain is particularly responsible for the directed localization of Intersectin to Clathrin-coated pits near the plasma membrane. Within the endocytic vesicles the SH3 domains facilitate the binding of Intersectin with Dynamin, and the central domain is essential for the association of Intersectin with SNAP 25. Two isoforms of Intersectin are produced as a result of alternative splicing in a stop codon, and they are designated as Intersectin-short and long (or Intersectin_S and Intersectin_L) to reflect an extended C-terminal domain. The long form, which has an extended C-terminal domain, is specifically expressed in neurons; the short form is detected in both glial and nonneuronal cells. The related proteins Intersectin-2 and the murine homolog Ese2 also contain the characteristic N-terminal EH domains, the central coiled-coil domain and five C-terminal SH3 domains and are likely involved the endocytic scaffolding complexes.

REFERENCES

1. Guipponi, M., Scott, H.S., Chen, H., Schebesta, A., Rossier, C. and Antonarakis, S.E. 1998. Two isoforms of a human Intersectin (ITSN) protein are produced by brain-specific alternative splicing in a stop codon. *Genomics* 53: 369-376.
2. Yamabhai, M., Hoffman, N.G., Hardison, N.L., McPherson, P.S., Castagnoli, L., Cesareni, G. and Kay, B.K. 1998. Intersectin, a novel adaptor protein with two Eps15 homology and five Src homology 3 domains. *J. Biol. Chem.* 273: 31401-31407.
3. Hussain, N.K., Yamabhai, M., Ramjaun, A.R., Guy, A.M., Baranes, D., O'Bryan, J.P., Der, C.J., Kay, B.K. and McPherson, P.S. 1999. Splice variants of Intersectin are components of the endocytic machinery in neurons and non-neuronal cells. *J. Biol. Chem.* 274: 15671-15677.

CHROMOSOMAL LOCATION

Genetic locus: ITSN2 (human) mapping to 2p23.3; Itsn2 (mouse) mapping to 12 A1.1.

SOURCE

Intersectin-2 (H-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Intersectin-2 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-9953 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Intersectin-2 (H-16) is recommended for detection of Intersectin-2_S and Intersectin-2_L of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Intersectin-2 (H-16) is also recommended for detection of Intersectin-2_S and Intersectin-2_L in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for Intersectin-2 siRNA (h): sc-41367, Intersectin-2 siRNA (m): sc-41368, Intersectin-2 shRNA Plasmid (h): sc-41367-SH, Intersectin-2 shRNA Plasmid (m): sc-41368-SH, Intersectin-2 shRNA (h) Lentiviral Particles: sc-41367-V and Intersectin-2 shRNA (m) Lentiviral Particles: sc-41368-V.

Positive Controls: Raji whole cell lysate: sc-364236, Jurkat whole cell lysate: sc-2204 or SUP-T1 whole cell lysate: sc-364796.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Intersectin-2 (H-16) : sc-9953. Western blot analysis of Intersectin-2 expression in Raji (A), Jurkat (B) and SUP-T1 (C) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.