MICAL3 (K-16): sc-99555



The Power to Question

BACKGROUND

MICAL3 (microtubule associated monoxygenase, calponin and LIM domain containing 3), also known as KIAA1364, is a 976 amino acid protein that localizes to both the cytoplasm and the cytoskeleton and contains one LIM zincbinding domain and one calponin-homology domain. Expressed ubiquitously, MICAL3 uses FAD as a cofactor to interact with Rab 1B and, via this interaction, is thought to play a role in spermatid development. MICAL3 exists as two alternatively spliced isoforms that are encoded by a gene which maps to human chromosome 22. Chromosome 22 houses over 500 genes, some of which are involved in Phelan-McDermid syndrome, schizophrenia and neurofibromatosis type 2.

REFERENCES

- Schwab, S.G. and Wildenauer, D.B. 1999. Chromosome 22 workshop report. Am. J. Med. Genet. 88: 276-278.
- Dunham, I., Shimizu, N., Roe, B.A., Chissoe, S., Hunt, A.R., Collins, J.E., Bruskiewich, R., Beare, D.M., Clamp, M., Smink, L.J., Ainscough, R., Almeida, J.P., Babbage, A., Bagguley, C., Bailey, J., Barlow, K., Bates, K.N., et al. 1999. The DNA sequence of human chromosome 22. Nature 402: 489-495.
- 3. Nagase, T., Kikuno, R., Ishikawa, K.I., Hirosawa, M. and Ohara, O. 2000. Prediction of the coding sequences of unidentified human genes. XVI. The complete sequences of 150 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 7: 65-73.
- Terman, J.R., Mao, T., Pasterkamp, R.J., Yu, H.H. and Kolodkin, A.L. 2002. MICALs, a family of conserved flavoprotein oxidoreductases, function in plexin-mediated axonal repulsion. Cell 109: 887-900.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608882. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/

CHROMOSOMAL LOCATION

Genetic locus: MICAL3 (human) mapping to 22q11.21; Mical3 (mouse) mapping to 6 F1.

SOURCE

MICAL3 (K-16) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of MICAL3 of human origin.

PRODUCT

Each vial contains 100 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-99555 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

MICAL3 (K-16) is recommended for detection of MICAL3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MICAL3 siRNA (h): sc-75783, MICAL3 siRNA (m): sc-149425, MICAL3 shRNA Plasmid (h): sc-75783-SH, MICAL3 shRNA Plasmid (m): sc-149425-SH, MICAL3 shRNA (h) Lentiviral Particles: sc-75783-V and MICAL3 shRNA (m) Lentiviral Particles: sc-149425-V.


Molecular Weight of MICAL3: 110 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, DU 145 cell lysate: sc-2268 or COLO 320DM cell lysate: sc-2226.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit lgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit lgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit lgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit lgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MICAL3 (K-16): sc-99555. Western blot analysis of MICAL3 expression in Hep G2 (**A**), Raji (**B**), DU 145 (**C**), SW480 (**D**) and COLO 320DM (**E**) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **MICAL3 (3A6): sc-517049**, our highly recommended monoclonal alternative to MICAL3 (K-16).