STAC (G-13): sc-99643



The Power to Question

BACKGROUND

STAC (SH3 and cysteine rich domain-containing protein), also known as STAC1, is a 402 amino acid protein that contains one SH3 (Src homology 3) domain and one cysteine-rich domain (CRD). Expressed in brain, STAC is a neuron-specific protein that localizes to the cytoplasm and, based on the frequent involvement of SH3 and CRD domains in signal transduction, is believed to play a role in neuron-specific signal transduction. In addition, STAC may be involved in protecting cells from apoptosis. Due to its neuron-specific expression and putative role in signal transduction, STAC may be implicated in a variety of hereditary neurological diseases.

REFERENCES

- Suzuki, H., Kawai, J., Taga, C., Yaoi, T., Hara, A., Hirose, K., Hayashizaki, Y. and Watanabe, S. 1996. STAC, a novel neuron-specific protein with cysteine-rich and SH3 domains. Biochem. Biophys. Res. Commun. 229: 902-909.
- Kawai, J., Suzuki, H., Hara, A., Hirose, K. and Watanabe, S. 1998. Human and mouse chromosomal mapping of STAC, a neuron-specific protein with an SH3 domain. Genomics 47: 140-142.
- Petek, E., Emberger, W., Kroisel, P.M. and Wagner, K. 1999. Assignment of STAC to human chromosome band 3p22.3 between D3S3718 and D3S1611. Cytogenet. Cell Genet. 84: 184-185.
- 4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 602317. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Hardy, K., Mansfield, L., Mackay, A., Benvenuti, S., Ismail, S., Arora, P., O'Hare, M.J. and Jat, P.S. 2005. Transcriptional networks and cellular senescence in human mammary fibroblasts. Mol. Biol. Cell 16: 943-953.
- Kato, Y., Uzawa, K., Saito, K., Nakashima, D., Kato, M., Nimura, Y., Seki, N. and Tanzawa, H. 2006. Gene expression pattern in oral cancer cervical lymph node metastasis. Oncol. Rep. 16: 1009-1014.
- 7. Satoh, J., Nanri, Y. and Yamamura, T. 2006. Rapid identification of 14-3-3-binding proteins by protein microarray analysis. J. Neurosci. Methods 152: 278-288.

CHROMOSOMAL LOCATION

Genetic locus: STAC (human) mapping to 3p22.3.

SOURCE

STAC (G-13) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of STAC of human origin.

PRODUCT

Each vial contains 100 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-99643 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

STAC (G-13) is recommended for detection of STAC of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with family members STAC2 or STAC3.

STAC (G-13) is also recommended for detection of STAC in additional species, including equine.

Suitable for use as control antibody for STAC siRNA (h): sc-78015, STAC shRNA Plasmid (h): sc-78015-SH and STAC shRNA (h) Lentiviral Particles: sc-78015-V.

Molecular Weight of STAC: 45 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com