

TROY (4E8): sc-100312



The Power to Question

BACKGROUND

The tumor necrosis factor (TNFR) superfamily represents a growing family of type I transmembrane glycoproteins that are involved in various cellular functions, including proliferation, differentiation and programmed cell death. These proteins share homology for cysteine-rich repeats in the extracellular ligand binding domain and an intracellular death domain. Members of the TNFR superfamily transmit signals through protein-protein interactions, and these signals can lead to the activation of either the caspase and Jun kinase pathways, which promote cell death, or the NF κ B pathway, which results in cell survival. One member of the TNFR superfamily TROY (also designated TAJ) exists as several isoforms, which vary in function. Full length TROY contains a cytoplasmic tail, which recruits tumor necrosis factor receptor-associated factor (TRAF) 2. The interaction between TROY and TRAF2 promotes cell survival through the NF κ B signaling pathway. TROY also exhibits significant homology to EDAR, a receptor that determines hair follicle fate, and like EDAR, TROY is expressed in the epithelium. Specifically, full length TROY mRNA is detected in the epithelium of mouse brain, embryo, heart, lung and liver. One truncated version of TROY, designated TNFRSF19, contains a shortened cytoplasmic tail, which prevents TNFRSF19 from activating the NF κ B signal transduction pathway.

REFERENCES

- Gruss, H.J. 1996. Molecular, structural, and biological characteristics of the tumor necrosis factor ligand superfamily. *Int. J. Clin. Lab. Res.* 26: 143-159.
- Gruss, H.J., et al. 1996. Structural and biological features of the TNF receptor and TNF ligand superfamilies: interactive signals in the pathobiology of Hodgkin's disease. *Ann. Oncol.* 7: 19-26.
- Baker, S.J., et al. 1998. Modulation of life and death by the TNF receptor superfamily. *Oncogene* 17: 3261-3270.
- Gurney, A.L., et al. 1999. Identification of a new member of the tumor necrosis factor family and its receptor, a human ortholog of mouse GITR. *Curr. Biol.* 9: 215-218.
- Hu, S., et al. 1999. Characterization of TNFRSF19, a novel member of the tumor necrosis factor receptor superfamily. *Genomics* 62: 103-107.
- Kojima, T., et al. 2000. TROY, a newly identified member of the tumor necrosis factor receptor superfamily, exhibits a homology with EDAR and is expressed in embryonic skin and hair follicles. *J. Biol. Chem.* 275: 20742-20747.
- Eby, M.T., et al. 2000. TAJ, a novel member of the tumor necrosis factor receptor family, activates the c-Jun N-terminal kinase pathway and mediates caspase-independent cell death. *J. Biol. Chem.* 275: 15336-15342.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: TNFRSF19 (human) mapping to 13q12.12.

SOURCE

TROY (4E8) is a mouse monoclonal antibody raised against recombinant TROY of human origin.

PRODUCT

Each vial contains 50 μ g IgG₁ kappa light chain in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

TROY (4E8) is recommended for detection of TROY of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TROY siRNA (h): sc-40247, TROY shRNA Plasmid (h): sc-40247-SH and TROY shRNA (h) Lentiviral Particles: sc-40247-V.

Molecular Weight of TROY: 45 kDa.


Positive Controls: LNCaP cell lysate: sc-2231 or A-431 whole cell lysate: sc-2201.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG_x BP-HRP: sc-516102 or m-IgG_x BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



TROY (4E8): sc-100312. Western blot analysis of TROY expression in HepG2 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.