

NDUFB3 (31-B): sc-100564

BACKGROUND

Complex 1 (also known as NADH dehydrogenase) of the electron transport chain (respiratory chain) is an enzymatic complex that catalyzes the transfer of electrons from NADH to ubiquinone. Free energy from the reaction is conserved in the transfer of protons into the intermembrane space to create an electrochemical proton gradient, a driving force for ATP synthesis. Complex 1 is a complicated, multi-protein, L-shaped complex composed of at least 45 different subunits and located in the mitochondrial inner membrane. NDUFB3 (NADH dehydrogenase (ubiquinone) 1 β subcomplex subunit 3), also known as B12, is a 98 amino acid hydrophobic protein belonging to the Complex I NDUFB3 subunit family. Ubiquitously expressed, NDUFB3 localizes to the matrix side of the inner membrane of the mitochondrion and functions as an accessory subunit of Complex I.

REFERENCES

1. Ton, C., et al. 1997. Identification and primary structure of five human NADH-ubiquinone oxidoreductase subunits. *Biochem. Biophys. Res. Commun.* 241: 589-594.
2. Loeffen, J.L., et al. 1998. cDNA of eight nuclear encoded subunits of NADH:ubiquinone oxidoreductase: human complex I cDNA characterization completed. *Biochem. Biophys. Res. Commun.* 253: 415-422.
3. Hadano, S., et al. 2001. Cloning and characterization of three novel genes, ALS2CR1, ALS2CR2, and ALS2CR3, in the juvenile amyotrophic lateral sclerosis (ALS2) critical region at chromosome 2q33-q34: candidate genes for ALS2. *Genomics* 71: 200-213.
4. Jensen, L.L., et al. 2001. Assignment of human NADH dehydrogenase (ubiquinone) 1 β subcomplex 3 (NDUFB3) and of its four pseudogenes to human chromosomes 2q31.3, 1p13.3→p13.1, 9q32→q34.1, 14q22.3→q23.1 and 14q32.2 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 93: 147-150.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603839. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Sparks, L.M., et al. 2005. A high-fat diet coordinately downregulates genes required for mitochondrial oxidative phosphorylation in skeletal muscle. *Diabetes* 54: 1926-1933.
7. Janssen, R.J., et al. 2006. Mitochondrial complex I: structure, function and pathology. *J. Inher. Metab. Dis.* 29: 499-515.
8. Edvardsen, H., et al. 2006. Experimental validation of data mined single nucleotide polymorphisms from several databases and consecutive dbSNP builds. *Pharmacogenet. Genomics* 16: 207-217.

CHROMOSOMAL LOCATION

Genetic locus: NDUFB3 (human) mapping to 2q33.1; Ndufb3 (mouse) mapping to 1 C1.3.

SOURCE

NDUFB3 (31-B) is a mouse monoclonal antibody raised against recombinant NDUFB3 of human origin.

PRODUCT

Each vial contains 100 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

NDUFB3 (31-B) is recommended for detection of NDUFB3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NDUFB3 siRNA (h): sc-94549, NDUFB3 siRNA (m): sc-149880, NDUFB3 shRNA Plasmid (h): sc-94549-SH, NDUFB3 shRNA Plasmid (m): sc-149880-SH, NDUFB3 shRNA (h) Lentiviral Particles: sc-94549-V and NDUFB3 shRNA (m) Lentiviral Particles: sc-149880-V.


Molecular Weight of NDUFB3: 12 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or A-431 whole cell lysate: sc-2201.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



NDUFB3 (31-B): sc-100564. Western blot analysis of NDUFB3 expression in HeLa (A) and A-431 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.