

UBE2H (18-Z): sc-100620

BACKGROUND

UBE2H (ubiquitin-conjugating enzyme E2H), also known as UBC8, UBCH, UBCH2 or E2-20K, is a 183 amino acid protein involved in ubiquitin-mediated protein degradation. Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). One of several members of the ubiquitin-conjugating enzyme family, UBE2H functions as an E2 ubiquitin-conjugating enzyme that acts to catalyze the covalent attachment of ubiquitin residues to various proteins, including Histone H2A. UBE2H shares 100% identity with its mouse counterpart and 98% identity with its frog and zebra-fish homologs, suggesting a conserved function between species. Multiple isoforms of UBE2H exist due to alternative splicing events.

REFERENCES

1. Kaiser, P., et al. 1994. A human ubiquitin-conjugating enzyme homologous to yeast UBC8. *J. Biol. Chem.* 269: 8797-8802.
2. Kaiser, P., et al. 1995. Characterization of functionally independent domains in the human ubiquitin conjugating enzyme UBCH2. *FEBS Lett.* 377: 193-196.
3. Wefes, I., et al. 1995. Induction of ubiquitin-conjugating enzymes during terminal erythroid differentiation. *Proc. Natl. Acad. Sci. USA* 92: 4982-4986.

CHROMOSOMAL LOCATION

Genetic locus: UBE2H (human) mapping to 7q32.2; Ube2h (mouse) mapping to 6 A3.3.

SOURCE

UBE2H (18-Z) is a mouse monoclonal antibody raised against recombinant UBE2H of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2a} kappa light chain in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

UBE2H (18-Z) is recommended for detection of UBE2H of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for UBE2H siRNA (h): sc-89802, UBE2H siRNA (m): sc-106659, UBE2H shRNA Plasmid (h): sc-89802-SH, UBE2H shRNA Plasmid (m): sc-106659-SH, UBE2H shRNA (h) Lentiviral Particles: sc-89802-V and UBE2H shRNA (m) Lentiviral Particles: sc-106659-V.


Molecular Weight of UBE2H: 21 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



UBE2H (18-Z): sc-100620. Western blot analysis of UBE2H expression in HeLa whole cell lysate.



UBE2H (18-Z): sc-100620. Immunofluorescence staining of paraformaldehyde-fixed HeLa cells showing nuclear localization.

SELECT PRODUCT CITATIONS

1. McKelvey, A.C., et al. 2016. RING finger E3 ligase PPP1R11 regulates TLR2 signaling and innate immunity. *Elife* 5: e18496.
2. Lim, K.H., et al. 2020. Predictive potential of circulating UBE2H mRNA as an E2 ubiquitin-conjugating enzyme for diagnosis or treatment of Alzheimer's disease. *Int. J. Mol. Sci.* 21: 3398.
3. Lear, T.B., et al. 2023. E3 ubiquitin ligase ZBTB25 suppresses β coronavirus infection through ubiquitination of the main viral protease MPro. *J. Biol. Chem.* 299: 105388.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.