

Act1 (WW-18): sc-100647

BACKGROUND

Members of the NF κ B family of transcription factors are important in regulating the expression of various cellular and viral genes involved in immune and inflammatory responses, cell survival and stress responses. IL-1, TNF α and other related signaling pathways activate transcription factors through the activation of JNK. The NF κ B signaling pathway converges with the signal-induced activation of JNK upstream of IKK. Isolated from the human embryonic kidney (HEK) 293 cell line, Act1 is an IKK γ -associated protein that activates both NF κ B and JNK constitutively. Act1, also designated NF κ B activator 1 or CIKS (for connection to IKK and SAPK/JNK), may function as a coordinator between two stress-induced signaling pathways.

REFERENCES

1. Siebenlist, U., et al. 1994. Structure, regulation and function of NF κ B. *Annu. Rev. Cell Biol.* 10: 405-455.
2. Barnes, P.J. and Karin, M. 1997. NF κ B: a pivotal transcription factor in chronic inflammatory diseases. *N. Engl. J. Med.* 336: 1066-1071.

CHROMOSOMAL LOCATION

Genetic locus: TRAF3IP2 (human) mapping to 6q21; Traf3ip2 (mouse) mapping to 10 B1.

SOURCE

Act1 (WW-18) is a mouse monoclonal antibody raised against recombinant Act1 of human origin.

PRODUCT

Each vial contains 100 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Act1 (WW-18) is recommended for detection of Act1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Act1 siRNA (h): sc-29634, Act1 siRNA (m): sc-29635, Act1 shRNA Plasmid (h): sc-29634-SH, Act1 shRNA Plasmid (m): sc-29635-SH, Act1 shRNA (h) Lentiviral Particles: sc-29634-V and Act1 shRNA (m) Lentiviral Particles: sc-29635-V.

Molecular Weight of Act1: 72 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227, Hep G2 nuclear extract: sc-364819 or HeLa nuclear extract: sc-2120.


STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



Act1 (WW-18): sc-100647. Western blot analysis of Act1 expression in Hep G2 nuclear extract.

Act1 (WW-18): sc-100647. Immunofluorescence staining of paraformaldehyde-fixed HeLa cells showing nuclear and cytoplasmic localization (A). Immunoperoxidase staining of formalin-fixed, paraffin-embedded human small intestine tissue showing membrane and cytoplasmic localization (B).

SELECT PRODUCT CITATIONS

1. Yariswamy, M., et al. 2016. Cardiac-restricted overexpression of TRAF3 interacting protein 2 (TRAF3IP2) results in spontaneous development of myocardial hypertrophy, fibrosis, and dysfunction. *J. Biol. Chem.* 291: 19425-19436.
2. Erikson, J.M., et al. 2017. Targeting TRAF3IP2 by genetic and interventional approaches inhibits ischemia/reperfusion-induced myocardial injury and adverse remodeling. *J. Biol. Chem.* 292: 2345-2358.
3. Sommer, A., et al. 2018. Th17 lymphocytes induce neuronal cell death in a human iPSC-based model of Parkinson's disease. *Cell Stem Cell* 23: 123-131.e6.
4. Abdellatif, A.M., et al. 2019. Human islet response to selected type 1 diabetes-associated bacteria: a transcriptome-based study. *Front. Immunol.* 10: 2623.
5. Ma, T., et al. 2020. Maternal exposure to di-n-butyl phthalate promotes the formation of testicular tight junctions through down-regulation of NF κ B/Cox-2/PGE2/MMP-2 in mouse offspring. *Environ. Sci. Technol.* 54: 8245-8258.
6. Xiao, D., et al. 2021. CRISPR-mediated rapid generation of neural cell-specific knockout mice facilitates research in neurophysiology and pathology. *Mol. Ther. Methods Clin. Dev.* 20: 755-764.
7. Pires, B.R.B., et al. 2021. Twist1 influences the expression of leading members of the IL-17 signaling pathway in HER2-positive breast cancer cells. *Int. J. Mol. Sci.* 22: 12144.
8. Miyashita, Y., et al. 2022. TICAM-1/TRIF associates with Act1 and suppresses IL-17 receptor-mediated inflammatory responses. *Life Sci. Alliance* 5: e202101181.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.