

MARCKS (JK-8): sc-100777

BACKGROUND

Myristoylated alanine-rich protein kinase C substrate (MARCKS), also designated 80K or 80K-L, has been identified as a major cellular substrate for protein kinase C. Human MARCKS is a 332 amino acid protein. The plasma membrane bound protein dissociates from the membrane upon phosphorylation by various PKC isoforms. In NIH/3T3 fibroblasts, PKC α and PKC ϵ , but not PKC δ , are responsible for MARCKS phosphorylation. MARCKS has been found to bind Calmodulin, Actin and Synapsin and is a filamentous (F) Actin crosslinking protein.

CHROMOSOMAL LOCATION

Genetic locus: MARCKS (human) mapping to 6q21; Marcks (mouse) mapping to 10 B1.

SOURCE

MARCKS (JK-8) is a mouse monoclonal antibody raised against amino acids 2-66 of MARCKS of human origin.

PRODUCT

Each vial contains 50 μ g IgG₁ kappa light chain in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

MARCKS (JK-8) is recommended for detection of MARCKS of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MARCKS siRNA (h): sc-35857, MARCKS siRNA (m): sc-35858, MARCKS shRNA Plasmid (h): sc-35857-SH, MARCKS shRNA Plasmid (m): sc-35858-SH, MARCKS shRNA (h) Lentiviral Particles: sc-35857-V and MARCKS shRNA (m) Lentiviral Particles: sc-35858-V.

Molecular Weight of MARCKS: 80 kDa.

Positive Controls: MARCKS (h2): 293T Lysate: sc-177518, SK-N-SH cell lysate: sc-2410 or NIH/3T3 whole cell lysate: sc-2210.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



MARCKS (JK-8): sc-100777. Western blot analysis of MARCKS expression in non-transfected 293T: sc-117752 (A), human MARCKS transfected 293T: sc-177518 (B) and SK-N-SH (C) whole cell lysates.



MARCKS (JK-8): sc-100777. Immunofluorescence staining of paraformaldehyde-fixed HeLa cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin-fixed, paraffin-embedded human spleen tissue showing cytoplasmic localization (B).

SELECT PRODUCT CITATIONS

- Shi, J., et al. 2014. Myristoylated alanine-rich C kinase substrate coordinates native TRPC1 channel activation by phosphatidylinositol 4,5-bisphosphate and protein kinase C in vascular smooth muscle. *FASEB J.* 28: 244-255.
- Fujita, K., et al. 2016. HMGB1, a pathogenic molecule that induces neurite degeneration via TLR4-MARCKS, is a potential therapeutic target for Alzheimer's disease. *Sci. Rep.* 6: 31895.
- Liu, H., et al. 2017. MiR-34c-3p acts as a tumor suppressor gene in osteosarcoma by targeting MARCKS. *Mol. Med. Rep.* 15: 1204-1210.
- Dao, C.V., et al. 2017. The MARCKS protein amount is differently regulated by Calpain during toxic effects of methylmercury between SH-SY5Y and EA.hy926 cells. *J. Vet. Med. Sci.* 79: 1931-1938.
- Wang, C.N., et al. 2019. Targeting the phosphorylation site of myristoylated alanine-rich C kinase substrate alleviates symptoms in a murine model of steroid-resistant asthma. *Br. J. Pharmacol.* 176: 1122-1134.
- Cilleros-Mañé, V., et al. 2021. M1 and M2 mAChRs activate PDK1 and regulate PKC β 1 and ϵ and the exocytotic apparatus at the NMJ. *FASEB J.* 35: e21724.
- Tanaka, H., et al. 2021. HMGB1 signaling phosphorylates Ku70 and impairs DNA damage repair in Alzheimer's disease pathology. *Commun. Biol.* 4: 1175.
- Long, X., et al. 2022. Obesity modulates cell-cell interactions during ovarian folliculogenesis. *iScience* 25: 103627.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.