

SH3BGR (46.1): sc-100853

BACKGROUND

Proline-rich peptide sequences have been shown to play important roles in protein-protein interactions that occur in signal transduction pathways. SH3 domain binding glutamic acid-rich protein (SH3BGR), also designated 21-glutamic acid-rich protein (21-GARP), is a 239 amino acid protein differentially expressed in heart and skeletal muscle. Its proline-rich region contains the consensus sequence for an SH3-binding domain and its acidic C-terminal region contains a glutamic acid-rich domain which may assume a coiled-coil structure. SH3BGR may be part of a multimeric complex, as it contains two functional domains involved in protein-protein interactions. The SH3BGR gene maps proximal to HMG14 on chromosome 21q22.3.

REFERENCES

- Scartezzini, P., Egeo, A., Colella, S., Fumagalli, P., Arrigo, P., Nizetic, D., Taramelli, R. and Rasore-Quartino, A. 1997. Cloning a new human gene from chromosome 21q22.3 encoding a glutamic acid-rich protein expressed in heart and skeletal muscle. *Hum. Genet.* 99: 387-392.
- Egeo, A., Mazzocco, M., Arrigo, P., Vidal-Taboada, J.M., Oliva, R., Pirola, B., Giglio, S., Rasore-Quartino, A. and Scartezzini, P. 1998. Identification and characterization of a new human gene encoding a small protein with high homology to the proline-rich region of the SH3BGR gene. *Biochem. Biophys. Res. Commun.* 247: 302-306.
- Mazzocco, M., Maffei, M., Egeo, A., Vergano, A., Arrigo, P., Di Lisi, R., Ghiotto, F. and Scartezzini, P. 2002. The identification of a novel human homologue of the SH3 binding glutamic acid-rich (SH3BGR) gene establishes a new family of highly conserved small proteins related to thioredoxin superfamily. *Gene* 291: 233-239.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 602230. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Majid, S.M., Liss, A.S., You, M. and Bose, H.R. 2005. The suppression of SH3BGR is important for v-Rel-mediated transformation. *Oncogene* 25: 756-768.
- Xu, C., Zheng, P., Shen, S., Xu, Y., Wei, L., Gao, H., Wang, S., Zhu, C., Tang, Y., Wu, J., Zhang, Q. and Shi, Y. 2005. NMR structure and regulated expression in APL cell of human SH3BGR3. *FEBS Lett.* 579: 2788-2794.

CHROMOSOMAL LOCATION

Genetic locus: SH3BGR (human) mapping to 21q22.2.

SOURCE

SH3BGR (46.1) is a mouse monoclonal antibody raised against recombinant SH3BGR of human origin.

PRODUCT

Each vial contains 100 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

SH3BGR (46.1) is recommended for detection of SH3BGR of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SH3BGR siRNA (h): sc-61543, SH3BGR shRNA Plasmid (h): sc-61543-SH and SH3BGR shRNA (h) Lentiviral Particles: sc-61543-V.


Molecular Weight of SH3BGR: 23 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



SH3BGR (46.1): sc-100853. Western blot analysis of SH3BGR expression in HeLa whole cell lysate.



SH3BGR (46.1): sc-100853. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic localization.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.