


WDR4 (321.9): sc-100894



BACKGROUND

WD-repeats are motifs that are found in a variety of proteins and are characterized by a conserved core of 40-60 amino acids that commonly form a tertiary propeller structure. While proteins that contain WD-repeats participate in a wide range of cellular functions, they are generally involved in regulatory mechanisms concerning chromatin assembly, cell cycle control, signal transduction, RNA processing, apoptosis and vesicular trafficking. WDR4 (WD-repeat-containing protein 4), also known as TRM82, is a 412 amino acid protein that contains two WD-repeats. Expressed as multiple isoforms due to alternative splicing events, WDR4 forms a complex with METTL1 (methyltransferase like 1) that is necessary for the 7-methylguanosine modification of tRNA. Defects in the gene encoding WDR4 may be associated with the development of Down syndrome, a chromosomal disorder characterized by deformed physical features and mental retardation.

REFERENCES

1. van der Voorn, L. and Ploegh, H.L. 1992. The WD-40 repeat. *FEBS Lett.* 307: 131-134.
2. Neer, E.J., et al. 1994. The ancient regulatory-protein family of WD-repeat proteins. *Nature* 371: 297-300.
3. Garcia-Higuera, I., et al. 1996. Folding of proteins with WD-repeats: comparison of six members of the WD-repeat superfamily to the G protein β subunit. *Biochemistry* 35: 13985-13994.

CHROMOSOMAL LOCATION

Genetic locus: WDR4 (human) mapping to 21q22.3.

SOURCE

WDR4 (321.9) is a mouse monoclonal antibody raised against recombinant WDR4 of human origin.

PRODUCT

Each vial contains 100 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

WDR4 (321.9) is recommended for detection of WDR4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for WDR4 siRNA (h): sc-91529, WDR4 shRNA Plasmid (h): sc-91529-SH and WDR4 shRNA (h) Lentiviral Particles: sc-91529-V.

Molecular Weight of WDR4: 45 kDa.


Positive Controls: HeLa nuclear extract: sc-2120, HeLa whole cell lysate: sc-2200 or A549 cell lysate: sc-2413.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG_κ BP-HRP: sc-516102 or m-IgG_κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG_κ BP-FITC: sc-516140 or m-IgG_κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.
- 4) Immunohistochemistry: use m-IgG_κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



WDR4 (321.9): sc-100894. Western blot analysis of WDR4 expression in HeLa nuclear extract.



WDR4 (321.9): sc-100894. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human small Intestine tissue showing nuclear localization.

SELECT PRODUCT CITATIONS

1. Zhang, L.S., et al. 2019. Transcriptome-wide mapping of internal N7-methylguanosine methylome in mammalian mRNA. *Mol. Cell* 74: 1304-1316.e8.
2. Zhao, Z., et al. 2023. QKI shuttles internal m⁷G-modified transcripts into stress granules and modulates mRNA metabolism. *Cell* 186: 3208-3226.e27.
3. Zhao, P., et al. 2024. METTL1 mediated tRNA m⁷G modification promotes leukaemogenesis of AML via tRNA regulated translational control. *Exp. Hematol. Oncol.* 13: 8.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.