

WDR20 (38K): sc-100900

BACKGROUND

WD-repeats are motifs that are found in a variety of proteins and are characterized by a conserved core of 40-60 amino acids that commonly form a tertiary propeller structure. While proteins that contain WD-repeats participate in a wide range of cellular functions, they are generally involved in regulatory mechanisms concerning chromatin assembly, cell cycle control, signal transduction, RNA processing, apoptosis and vesicular trafficking. WDR20 (WD repeat-containing protein 20), also known as DMR, is a 569 amino acid protein that contains 5 WD-repeats and may be involved in signaling networks throughout the cell. Due to alternative splicing events, two isoforms of WDR20 are expressed.

REFERENCES

1. van der Voorn, L. and Ploegh, H.L. 1992. The WD-40 repeat. *FEBS Lett.* 307: 131-134.
2. Neer, E.J., et al. 1994. The ancient regulatory-protein family of WD-repeat proteins. *Nature* 371: 297-300.
3. Garcia-Higuera, I., et al. 1996. Folding of proteins with WD-repeats: comparison of six members of the WD-repeat superfamily to the G protein β subunit. *Biochemistry* 35: 13985-13994.

CHROMOSOMAL LOCATION

Genetic locus: WDR20 (human) mapping to 14q32.31; Wdr20 (mouse) mapping to 12 F1.

SOURCE

WDR20 (38K) is a mouse monoclonal antibody raised against recombinant WDR20 of human origin.

PRODUCT

Each vial contains 100 μ g IgG₁ kappa light chain in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

WDR20 (38K) is recommended for detection of WDR20 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for WDR20 siRNA (h): sc-92396, WDR20 shRNA Plasmid (h): sc-92396-SH and WDR20 shRNA (h) Lentiviral Particles: sc-92396-V.


Molecular Weight of WDR20: 63 kDa.

Positive Controls: WDR20 (h): 293T Lysate: sc-174532 or HeLa nuclear extract: sc-2120.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



WDR20 (38K): sc-100900. Western blot analysis of WDR20 expression in non-transfected: sc-117752 (A) and human WDR20 transfected: sc-174532 (B) 293T whole cell lysates and HeLa nuclear extract (C).



WDR20 (38K): sc-100900. Immunofluorescence staining of paraformaldehyde-fixed HeLa cells showing nuclear localization (A). Immunoperoxidase staining of formalin-fixed, paraffin-embedded human colon tissue showing nuclear localization (B).

SELECT PRODUCT CITATIONS

1. McClurg, U.L., et al. 2015. Ubiquitin-specific protease 12 interacting partners Uaf-1 and WDR20 are potential therapeutic targets in prostate cancer. *Oncotarget* 6: 37724-37736.
2. Ju, L.G., et al. 2018. Characterization of WDR20: a new regulator of the ERAD machinery. *Biochim. Biophys. Acta* 1865: 970-980.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.