

SH3RF2 (Y-18): sc-100976

BACKGROUND

The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. SH3RF2 (SH3 domain containing RING finger 2), also known as RNF158, is a 729 amino acid protein with one RING-type zinc finger domain and 3 SH3 domains. Via its RING-type zinc finger domain, SH3RF2 binds an E2 ubiquitin-conjugating enzyme. This suggests that SH3RF2 functions as an E3 ubiquitin-protein ligase that accepts a ubiquitin residue from an E2 ubiquitin-conjugating enzyme and immediately transfers that residue to a protein that is targeted for degradation. Due to alternative splicing events, SH3RF2 is expressed as two different isoforms.

REFERENCES

- Gerhard, D.S., Wagner, L., Feingold, E.A., Shenmen, C.M., Grouse, L.H., Schuler, G., Klein, S.L., Old, S., Rasooly, R., Good, P., Guyer, M., Peck, A.M., Derge, J.G., Lipman, D., Collins, F.S., Jang, W., Sherry, S., et al. 2004. The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). *Genome Res.* 14: 2121-2127.
- Ota, T., Suzuki, Y., Nishikawa, T., Otsuki, T., Sugiyama, T., Irie, R., Wakamatsu, A., Hayashi, K., Sato, H., Nagai, K., Kimura, K., Makita, H., Sekine, M., Obayashi, M., Nishi, T., Shibahara, T., Tanaka, T., et al. 2004. Complete sequencing and characterization of 21,243 full-length human cDNAs. *Nat. Genet.* 36: 40-45.
- Ernst, C., Sequeira, A., Klempan, T., Ernst, N., French-Mullen, J. and Turecki, G. 2007. Confirmation of region-specific patterns of gene expression in the human brain. *Neurogenetics* 8: 219-224.

CHROMOSOMAL LOCATION

Genetic locus: SH3RF2 (human) mapping to 5q32; Sh3rf2 (mouse) mapping to 18 B3.

SOURCE

SH3RF2 (Y-18) is a mouse monoclonal antibody raised against recombinant SH3RF2 of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

SH3RF2 (Y-18) is recommended for detection of SH3RF2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for SH3RF2 siRNA (h): sc-91808, SH3RF2 siRNA (m): sc-153440, SH3RF2 shRNA Plasmid (h): sc-91808-SH, SH3RF2 shRNA Plasmid (m): sc-153440-SH, SH3RF2 shRNA (h) Lentiviral Particles: sc-91808-V and SH3RF2 shRNA (m) Lentiviral Particles: sc-153440-V.


Molecular Weight of SH3RF2: 79 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or HeLa nuclear extract: sc-2120.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



SH3RF2 (Y-18): sc-100976. Western blot analysis of SH3RF2 expression in HeLa whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.