

HNF-4 α (Q-15): sc-101059

BACKGROUND

HNF-1 (α and β), HNF-3 (α , β and γ), HNF-4 (α and γ) and HNF-6 compose, in part, a homeoprotein family designated the hepatocyte nuclear factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. HNF-3 α , HNF-3 β and HNF-3 γ regulate the transcription of numerous hepatocyte genes in adult liver. HNF-3 α and HNF-3 β have also been shown to be involved in gastrulation events such as body axis formation. HNF-4 α and HNF-4 γ have been shown to be important for early embryo development. HNF-4 α is expressed in liver, kidney, pancreas, small intestine, testis and colon; HNF-4 γ is expressed in each of these tissues except liver. HNF-6 has been shown to bind to the promoter of HNF-3 β , which indicates a potential role of HNF-6 in gut endoderm epithelial cell differentiation. Evidence suggests that HNF-6 may also be a transcriptional activator for at least 22 other hepatocyte-enriched genes, including cytochrome P450 2C13 and α -1 antitrypsin.

REFERENCES

1. Bach, I., et al. 1993. More potent transcriptional activators or a trans-dominant inhibitor of the HNF1 homeoprotein family are generated by alternative RNA processing. *EMBO J.* 12: 4229-4242.
2. Kaestner, K.H., et al. 1994. The HNF-3 gene family of transcription factors in mice: gene structure, cDNA sequence, and mRNA distribution. *Genomics* 20: 377-385.

CHROMOSOMAL LOCATION

Genetic locus: HNF4A (human) mapping to 20q13.12; Hnf4a (mouse) mapping to 2 H3.

SOURCE

HNF-4 α (Q-15) is a mouse monoclonal antibody raised against recombinant HNF-4 α of human origin.

PRODUCT

Each vial contains 50 μ g IgG_{2a} kappa light chain in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

HNF-4 α (Q-15) is recommended for detection of HNF-4 α of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HNF-4 α siRNA (h): sc-35573, HNF-4 α siRNA (m): sc-35574, HNF-4 α shRNA Plasmid (h): sc-35573-SH, HNF-4 α shRNA Plasmid (m): sc-35574-SH, HNF-4 α shRNA (h) Lentiviral Particles: sc-35573-V and HNF-4 α shRNA (m) Lentiviral Particles: sc-35574-V.


Molecular Weight of full-length HNF-4 α : 54 kDa.

Molecular Weight of N-terminal truncated HNF-4 α : 40 kDa.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



HNF-4 α (Q-15): sc-101059. Western blot analysis of HNF-4 α expression in Hep G2 whole cell lysate.



HNF-4 α (Q-15): sc-101059. Immunofluorescence staining of paraformaldehyde-fixed HeLa cells showing nuclear localization.

SELECT PRODUCT CITATIONS

1. Felder, T.K., et al. 2011. Characterization of novel peroxisome proliferator-activated receptor γ coactivator-1 α (PGC-1 α) isoform in human liver. *J. Biol. Chem.* 286: 42923-42936.
2. Wang, G., et al. 2018. Co-culture system of hepatocytes and endothelial cells: two *in vitro* approaches for enhancing liver-specific functions of hepatocytes. *Cytotechnology* 70: 1279-1290.
3. Ge, W., et al. 2023. Circadian PER1 controls daily fat absorption with the regulation of PER1-PKA on phosphorylation of bile acid synthetase. *J. Lipid Res.* 64: 100390.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.