

PSIP1 (3F7): sc-101087

BACKGROUND

PSIP1 (PC4 and SFRS1 interacting protein 1), also known as CLL-associated antigen KW-7, PSIP2, LEDGF (lens epithelium-derived growth factor), PAIP, DFS70 (dense fine speckles 70 kDa protein) or transcriptional coactivator p75/p52, is a 530 amino acid nuclear protein that associates with chromatin throughout the cell cycle. Functioning as a transcriptional coactivator that complexes with the human immunodeficiency virus type 1 (HIV-1) integrase, PSIP1 is essential for the nuclear localization and chromosomal association of viral proteins. As the primary integrase-to-chromatin tethering factor for HIV-1, PSIP1 is responsible for the cellular trafficking of lentiviral integrases. During apoptosis, PSIP1 is cleaved at three sites by caspase-3 and caspase-7, contributing to the pathogenesis of atopic disorders.

REFERENCES

- Ge, H., et al. 1998. Isolation of cDNAs encoding novel transcription coactivators p52 and p75 reveals an alternate regulatory mechanism of transcriptional activation. *EMBO J.* 17: 6723-6729.
- Singh, D.P., et al. 2000. Lens epithelium-derived growth factor (LEDGF/p75) and p52 are derived from a single gene by alternative splicing. *Gene* 242: 265-273.
- Krackhardt, A.M., et al. 2002. Identification of tumor-associated antigens in chronic lymphocytic leukemia by SEREX. *Blood* 100: 2123-2131.
- Wu, X., et al. 2002. Caspase cleavage of the nuclear autoantigen LEDGF/p75 abrogates its pro-survival function: implications for autoimmunity in atopic disorders. *Cell Death Differ.* 9: 915-925.
- Cherepanov, P., et al. 2003. HIV-1 integrase forms stable tetramers and associates with LEDGF/p75 protein in human cells. *J. Biol. Chem.* 278: 372-381.
- Cherepanov, P., et al. 2004. Identification of an evolutionarily conserved domain in human lens epithelium-derived growth factor/transcriptional co-activator p75 (LEDGF/p75) that binds HIV-1 integrase. *J. Biol. Chem.* 279: 48883-48892.

CHROMOSOMAL LOCATION

Genetic locus: PSIP1 (human) mapping to 9p22.3; Psip1 (mouse) mapping to 4 C3.

SOURCE

PSIP1 (3F7) is a mouse monoclonal antibody raised against recombinant PSIP1 of human origin.

PRODUCT

Each vial contains 100 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

PSIP1 (3F7) is recommended for detection of PSIP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PSIP1 siRNA (h): sc-44991, PSIP1 siRNA (m): sc-44992, PSIP1 shRNA Plasmid (h): sc-44991-SH, PSIP1 shRNA Plasmid (m): sc-44992-SH, PSIP1 shRNA (h) Lentiviral Particles: sc-44991-V and PSIP1 shRNA (m) Lentiviral Particles: sc-44992-V.

Molecular Weight of PSIP1 p52 isoform: 38 kDa.


Molecular Weight of PSIP1 p75 isoform: 60 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



PSIP1 (3F7): sc-101087. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human testis tissue showing nuclear localization.

SELECT PRODUCT CITATIONS

- Knyazhanskaya, E., et al. 2019. NHEJ pathway is involved in post-integration DNA repair due to Ku70 binding to HIV-1 integrase. *Retrovirology* 16: 30.

RESEARCH USE

For research use only, not for use in diagnostic procedures.