# TLE5 (F-G9): sc-101181



The Power to Question

### **BACKGROUND**

The Notch signaling pathway controls cellular interactions important for the specification of a variety of fates in both vertebrates and invertebrates. Key players in the Notch pathway are the TLE genes (for transducin-like enhancer of split, also designated ESG for enhancer of split groucho), which are human homologs of the *Drosophila* groucho gene. TLE5, also known as AES (aminoterminal enhancer of split), GRG or ESP1, is a 197 amino acid nuclear protein that belongs to the TLE family. Expressed predominately in fetal brain, liver, lung, heart and kidney and in adult muscle, TLE5 functions as either a homoligomer or a heterooligomer with other TLE family members and, through this association, dominantly represses the expression of TLE genes. In addition, TLE5 can repress  $NF\kappa B$ -regulated gene expression and is thought to play an important role in initiating and maintaining cell differentiation events. Two isoforms of TLE5 exist due to alternative splicing events.

## **REFERENCES**

- Miyasaka, H., et al. 1993. Molecular cloning and expression of mouse and human cDNA encoding AES and ESG proteins with strong similarity to *Drosophila* enhancer of split groucho protein. Eur. J. Biochem. 216: 343-352.
- Hou, E.W. and Li, S.S. 1998. Genomic organization and chromosome localization to band 19p13.3 of the human AES gene: gene product exhibits strong similarity to the N-terminal domain of *Drosophila* enhancer of split groucho protein. DNA Cell Biol. 17: 911-913.
- 3. Tetsuka, T., et al. 2000. Inhibition of NF $\kappa$ B-mediated transcription by association with the amino-terminal enhancer of split, a groucho-related protein lacking WD40 repeats. J. Biol. Chem. 275: 4383-4390.
- Wang, J.C., et al. 2000. Transducin-like enhancer of split proteins, the human homologs of *Drosophila* groucho, interact with hepatic nuclear factor 3β. J. Biol. Chem. 275: 18418-18423.
- Yochum, G.S. and Ayer, D.E. 2001. PF-1, a novel PHD zinc finger protein that links the TLE corepressor to the mSin3A-histone deacetylase complex. Mol. Cell. Biol. 21: 4110-4118.
- López-Ríos, J., et al. 2003. Six3 and Six6 activity is modulated by members of the groucho family. Development 130: 185-195.
- Jan, Y., et al. 2004. A mitochondrial protein, Bit1, mediates apoptosis regulated by integrins and groucho/TLE corepressors. Cell 116: 751-762.
- 8. Sekiya, T. and Zaret, K.S. 2007. Repression by groucho/TLE/Grg proteins: genomic site recruitment generates compacted chromatin *in vitro* and impairs activator binding *in vivo*. Mol. Cell 28: 291-303.
- Jennings, B.H. and Ish-Horowicz, D. 2008. The groucho/TLE/Grg family of transcriptional co-repressors. Genome Biol. 9: 205.

# **CHROMOSOMAL LOCATION**

Genetic locus: AES (human) mapping to 19p13.3; Aes (mouse) mapping to 10 C1.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **SOURCE**

TLE5 (F-G9) is a mouse monoclonal antibody raised against recombinant TLE5 of human origin.

### **PRODUCT**

Each vial contains 100  $\mu g \ lgG_1$  kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### **APPLICATIONS**

TLE5 (F-G9) is recommended for detection of TLE5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TLE5 siRNA (h): sc-63133, TLE5 siRNA (m): sc-63134, TLE5 shRNA Plasmid (h): sc-63133-SH, TLE5 shRNA Plasmid (m): sc-63134-SH, TLE5 shRNA (h) Lentiviral Particles: sc-63133-V and TLE5 shRNA (m) Lentiviral Particles: sc-63134-V.


Molecular Weight of TLE5: 22 kDa.

Positive Controls: HeLa nuclear extract: sc-2120.

# **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG $\kappa$  BP-HRP: sc-516102 or m-lgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

#### **DATA**



TLE5 (F-G9): sc-101181. Western blot analysis of TLE5 expression in HeLa nuclear extract.

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.