

MAGE-A11 (YN-2): sc-101222

BACKGROUND

The melanoma-associated antigen (MAGE) family consists of a number of antigens recognized by cytotoxic T lymphocytes. The MAGE genes were initially isolated from different kinds of tumors and, based on their virtually exclusive tumor-specific expression in adult tissues, they have been used as targets for cancer immunotherapy. MAGE genes encode for tumor-rejection antigens and are expressed in tumors of different histologic types and in normal testis and placenta. MAGE-A11 (melanoma antigen family A, 11), also known as MAGE11, MAGE-11, MAGEA-11 or CT1.11 (cancer/testis antigen 1.11), is a 429 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one MAGE domain. Expressed in a variety of tumors, including melanoma, breast cancer and lung cancer, MAGE-A11 functions as an androgen receptor (AR) co-regulator that modulates the interdomain of AR, thereby increasing its activity. Through its regulation of AR, MAGE-A11 is thought to play an important role in embryonic development and tumor progression/transformation. Two isoforms of MAGE-A11 exist due to alternative splicing events.

REFERENCES

1. De Plaen, E., Arden, K., Traversari, C., Gaforio, J.J., Szikora, J.P., De Smet, C., Brasseur, F., van der Bruggen, P., Lethé, B. and Lurquin, C. 1994. Structure, chromosomal localization, and expression of 12 genes of the MAGE family. *Immunogenetics* 40: 360-369.
2. Rogner, U.C., Wilke, K., Steck, E., Korn, B. and Poustka, A. 1995. The melanoma antigen gene (MAGE) family is clustered in the chromosomal band Xq28. *Genomics* 29: 725-731.
3. Jurk, M., Kremmer, E., Schwarz, U., Förster, R. and Winnacker, E.L. 1998. MAGE-11 protein is highly conserved in higher organisms and located predominantly in the nucleus. *Int. J. Cancer* 75: 762-766.
4. Irvine, R.A. and Coetzee, G.A. 1999. Additional upstream coding sequences of MAGE-11. *Immunogenetics* 49: 585.
5. Serrano, A., Lethé, B., Delroisse, J.M., Lurquin, C., De Plaen, E., Brasseur, F., Rimoldi, D. and Boon, T. 1999. Quantitative evaluation of the expression of MAGE genes in tumors by limiting dilution of cDNA libraries. *Int. J. Cancer* 83: 664-669.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 300344: World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Bai, S., He, B. and Wilson, E.M. 2005. Melanoma antigen gene protein MAGE-11 regulates androgen receptor function by modulating the inter-domain interaction. *Mol. Cell. Biol.* 25: 1238-1257.
8. Askew, E.B., Gampe, R.T., Stanley, T.B., Faggart, J.L. and Wilson, E.M. 2007. Modulation of androgen receptor activation function 2 by testosterone and dihydrotestosterone. *J. Biol. Chem.* 282: 25801-25816.
9. Bai, S. and Wilson, E.M. 2008. Epidermal-growth-factor-dependent phosphorylation and ubiquitinylation of MAGE-11 regulates its interaction with the androgen receptor. *Mol. Cell. Biol.* 28: 1947-1963.

CHROMOSOMAL LOCATION

Genetic locus: MAGEA11 (human) mapping to Xq28.

SOURCE

MAGE-A11 (YN-2) is a mouse monoclonal antibody raised against recombinant MAGE-A11 of human origin.

PRODUCT

Each vial contains 200 µl ascites containing IgM with < 0.1% sodium azide.

APPLICATIONS

MAGE-A11 (YN-2) is recommended for detection of MAGE-A11 of human origin by Western Blotting (starting dilution to be determined by researcher, dilution range 1:100-1:5000), immunoprecipitation [1-2 µl per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution to be determined by researcher, dilution range 1:100-1:5000).

Suitable for use as control antibody for MAGE-A11 siRNA (h): sc-108016, MAGE-A11 shRNA Plasmid (h): sc-108016-SH and MAGE-A11 shRNA (h) Lentiviral Particles: sc-108016-V.

Molecular Weight of full length MAGE-A11: 70 kDa.

Molecular Weight of truncated MAGE-A11: 40 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203.

STORAGE

For immediate and continuous use, store at 4° C for up to one month. For sporadic use, freeze in working aliquots in order to avoid repeated freeze/thaw cycles. If turbidity is evident upon prolonged storage, clarify solution by centrifugation.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.