

## IL-1ra (5G11): sc-101342

### BACKGROUND

Two forms of interleukin-1, designated IL-1 $\alpha$  and IL-1 $\beta$ , have been described. Although encoded by distinct genes and exhibiting roughly only 25% sequence identity, IL-1 $\alpha$  and IL-1 $\beta$  bind to the same receptor and seem to elicit similar biological responses. IL-1 production is generally thought to be associated with inflammation, but it has also been shown to be expressed during kidney development, thymocyte differentiation and cartilage degradation. IL-1 plays a critical role in the regulation of immune response and inflammation acting as an activator of T and B lymphocytes and natural killer (NK) cells. IL-1 receptor antagonist (IL-1ra) is a cytokine that inhibits IL-1 $\alpha$  and IL-1 $\beta$  binding to interleukin receptors. By neutralizing the activity of IL-1, IL-1ra contributes to the inhibition of the immune and inflammatory responses and has been targeted as a drug for the treatment of severely active rheumatoid arthritis. There are four isoforms of IL-1ra that are produced as a result of alternative splicing events.

### REFERENCES

1. Auron, P.E., et al. 1984. Nucleotide sequence of human monocyte interleukin-1 precursor cDNA. Proc. Natl. Acad. Sci. USA 81: 7907-7911.
2. March, C.J., et al. 1985. Cloning, sequence and expression of two distinct human interleukin-1 complementary DNAs. Nature 315: 641-647.
3. Carter, D.B., et al. 1990. Purification, cloning, expression and biological characterization of an interleukin-1 receptor antagonist protein. Nature 344: 633-638.
4. Sadouk, M.B., et al. 1995. Human synovial fibroblasts coexpress IL-1 receptor type I and type II mRNA. The increased level of the IL-1 receptor in osteoarthritic cells is related to an increased level of the type I receptor. Lab. Invest. 73: 347-355.
5. Lonnemann, G., et al. 1995. Cytokines in human renal interstitial fibrosis. I. Interleukin-1 is a paracrine growth factor for cultured fibrosis-derived kidney fibroblasts. Kidney Int. 47: 837-844.
6. Zuniga-Pflucker, J.C., et al. 1995. Requirement for TNF $\alpha$  and IL-1 $\alpha$  in fetal thymocyte commitment and differentiation. Science 268: 1906-1909.
7. Sandborg, C.I., et al. 1995. IL-4 expression in human T cells is selectively inhibited by IL-1 $\alpha$  and IL-1 $\beta$ . J. Immunol. 155: 5206-5212.

### CHROMOSOMAL LOCATION

Genetic locus: IL1RN (human) mapping to 2q13.

### SOURCE

IL-1ra (5G11) is a mouse monoclonal antibody raised against recombinant IL-1ra of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### APPLICATIONS

IL-1ra (5G11) is recommended for detection of IL-1ra of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500); non cross-reactive with IL-1 $\alpha$  or IL-1 $\beta$ .

Suitable for use as control antibody for IL-1ra siRNA (h): sc-39617; and as shRNA Plasmid control antibody for IL-1ra shRNA Plasmid (h): sc-39617-SH.

Molecular Weight of IL-1ra: 17-25 kDa.

### RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG $\kappa$  BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.