

GDF-15 (ME-6D10): sc-101379

BACKGROUND

Growth differentiation factor 15 (GDF-15), also known as PDF, MIC-1, PLAB, NAG-1 or PTGF- β , is a member of the transforming growth factor β (TGF- β) superfamily. Synthesized intracellularly, the protein is secreted as a dimer linked by disulfide bonds. Epithelial cells and macrophages are the sites of strongest GDF-15 expression, although it is widely expressed in adult tissue. In the brain, GDF-15 expression occurs in the choroid plexus, from which the protein is secreted into the cerebrospinal fluid. The gene for GDF-15 is responsive to p53 tumor suppressor protein, and in cultured cerebellar granule neurons GDF-15 can prevent cell death by the activation of Akt and inhibition of ERK. GDF-15 acts as a trophic factor for certain classes of neurons, promoting cell survival and differentiation. Overexpression of GDF-15 occurs in prostate cancer, and may be a means of diagnosis. In the uterus, GDF-15 may work to suppress maternally derived proinflammatory cytokines, thereby promoting fetal survival.

REFERENCES

1. Fairlie, W.D., et al. 1999. MIC1 is a novel TGF β superfamily cytokine associated with macrophage activation. *J. Leukoc. Biol.* 65: 2-5.
2. Bottner, M., et al. 1999. Expression of a novel member of the TGF β superfamily, growth/differentiation factor-15/macrophage-inhibiting cytokine-1 (GDF-15/MIC1) in adult rat tissues. *Cell Tissue Res.* 297: 103-110.
3. Strelau, J., et al. 2000. GDF-15/MIC1 a novel member of the TGF β superfamily. *J. Neural Transm. Suppl.* 60: 273-276.
4. Moore, A.G., et al. 2000. The transforming growth factor β superfamily cytokine macrophage inhibitory cytokine-1 is present in high concentrations in the serum of pregnant women. *J. Clin. Endocrinol. Metab.* 85: 4781-4788.
5. Bauskin, A.R., et al. 2000. The propeptide of macrophage inhibitory cytokine (MIC1), a TGF β superfamily member, acts as a quality control determinant for correctly folded MIC1. *EMBO J.* 19: 2212-2220.
6. Kim, K.S., et al. 2002. Expression and regulation of nonsteroidal anti-inflammatory drug-activated gene (NAG1) in human and mouse tissue. *Gastroenterology* 122: 1388-1398.
7. Kriegstein, K., et al. 2002. TGF β and the regulation of neuron survival and death. *J. Physiol. Paris* 96: 25-30.
8. Strelau, J., et al. 2003. Growth/differentiation factor-15 (GDF-15), a novel member of the TGF β superfamily, promotes survival of lesioned mesencephalic dopaminergic neurons *in vitro* and *in vivo* and is induced in neurons following cortical lesioning. *J. Neural Transm. Suppl.* 65: 197-203.
9. Iczkowski, K.A., et al. 2003. Overexpression of NSAID-activated gene product in prostate cancer. *Int. J. Surg. Pathol.* 11: 159-166.

CHROMOSOMAL LOCATION

Genetic locus: GDF15 (human) mapping to 19p13.11.

SOURCE

GDF-15 (ME-6D10) is a mouse monoclonal antibody genetically immunized with cDNA encoding amino acids 30-194 of GDF-15 of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

GDF-15 (ME-6D10) is available conjugated to either phycoerythrin (sc-101379 PE) or fluorescein (sc-101379 FITC), 200 μ g/ml, for IF, IHC(P) and FCM.

APPLICATIONS

GDF-15 (ME-6D10) is recommended for detection of GDF-15 of human origin by flow cytometry (1 μ g per 1 x 10⁶ cells) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GDF-15 siRNA (h): sc-39798, GDF-15 shRNA Plasmid (h): sc-39798-SH and GDF-15 shRNA (h) Lentiviral Particles: sc-39798-V.

Molecular Weight of GDF-15 precursor: 40 kDa.

Molecular Weight of mature GDF-15: 30 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.