

survivin (3C5B2): sc-101432

BACKGROUND

The baculovirus protein p35 inhibits virally-induced apoptosis of invertebrate and mammalian cells and may function to impair the clearing of virally-infected cells by the immune system of the host. This is accomplished, at least in part, by the ability of p35 to block both TNF- and FAS-mediated apoptosis through the inhibition of the ICE family of serine proteases. Two mammalian homologs of baculovirus p35, referred to as inhibitor of apoptosis protein (IAP) 1 and 2, share an amino-terminal baculovirus IAP repeat (BIR) motif and a carboxy-terminal RING finger. Although the c-IAPs do not directly associate with the TNF receptor (TNF-R), they efficiently block TNF-mediated apoptosis through their interaction with the downstream TNF-R effectors, TRAF1 and TRAF2. Additional IAP family members include ILP (for IAP-like protein) and survivin. ILP inhibits activated caspase-3, leading to the resistance of FAS-mediated apoptosis. Survivin (also designated TIAP) is expressed during the G₂/M phase of the cell cycle and associates with microtubules of the mitotic spindle. Increased caspase-3 activity is detected when a disruption of survivin-microtubule interactions occurs.

REFERENCES

- Hay, B.A., et al. 1994. Expression of baculovirus p35 prevents cell death in *Drosophila*. *Development* 120: 2121-2129.
- Beidler, D.R., et al. 1995. The baculovirus p35 protein inhibits FAS- and tumor necrosis factor-induced apoptosis. *J. Biol. Chem.* 270: 16526-16528.
- Bump, N.J., et al. 1995. Inhibition of ICE family proteases by baculovirus antiapoptotic protein p35. *Science* 269: 1885-1888.
- Rothe, M., et al. 1995. The TNF-R2-TRAF signaling complex contains two novel proteins related to baculoviral inhibitor of apoptosis proteins. *Cell* 83: 1243-1252.
- Uren, A.G., et al. 1996. Cloning and expression of apoptosis inhibitory protein homologs that function to inhibit apoptosis and/or bind tumor necrosis factor receptor-associated factors. *Proc. Natl. Acad. Sci. USA* 93: 4974-4978.
- Suzuki, A., et al. 1998. Resistance to Fas-mediated apoptosis: activation of caspase 3 is regulated by cell cycle regulator p21^{WAF1} and IAP gene family ILP. *Oncogene* 17: 931-939.
- Li, F., et al. 1998. Control of apoptosis and mitotic spindle checkpoint by survivin. *Nature* 396: 580-584.
- Ma, J. et al. 2007. Effects of Genistein on the proliferation and expression of survivin in salivary adenoid cystic carcinoma cell line SACC-83. *Hua Xi Kou Qiang Yi Xue Za Zhi* 25: 97-99.

CHROMOSOMAL LOCATION

Genetic locus: BIRC5 (human) mapping to 17q25.3.

SOURCE

survivin (3C5B2) is a mouse monoclonal antibody raised against recombinant full length survivin of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

survivin (3C5B2) is recommended for detection of survivin of human origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for survivin siRNA (h): sc-29499, survivin shRNA Plasmid (h): sc-29499-SH and survivin shRNA (h) Lentiviral Particles: sc-29499-V.

Molecular Weight of survivin: 17 kDa.

Positive Controls: COLO 320DM cell lysate: sc-2226, HL-60 whole cell lysate: sc-2209 or MOLT-4 cell lysate: sc-2233.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CONJUGATES

See **survivin (D-8): sc-17779** for survivin antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.