# SANTA CRUZ BIOTECHNOLOGY, INC.

# HSP 90 (AC-16): sc-101494



## BACKGROUND

The heat shock response was first described for *Drosophila* salivary gland cells and morphologically consists of a change in their polytene chromosome puffing patterns that involves *de novo* synthesis of a few proteins. Similar heat shock proteins were later discovered in bacterial chicken and mammalian cells, and have been subsequently studied in other organisms. A series of proteins, including HSP 90, HSP 70, HSP 20-30 and ubiquitin, are induced by insults such as temperature shock, chemicals and other environmental stress. A major function of HSP 90 and other HSPs is to act as molecular chaperones. HSP 90 forms a complex with glucocorticoid receptor (GR), rendering the non ligand-bound receptor transcriptionally inactive. HSP 90 binds the GR as a heterocomplex composed of either HSP 56 or Cyclophilin D, forming an aporeceptor complex. HSP 90 also exists as a dimer with other proteins such as p60/STI1 and p23, forming an apo-receptor complex with estrogen and androgen receptors.

# REFERENCES

- 1. Wu, J.M., et al. 2003. PKC  $\epsilon$  is a unique regulator for HSP 90 $\beta$  gene in heat shock response. J. Biol. Chem. 278: 51143-51149.
- 2. Whitesell, L., et al. 2005. HSP 90 and the chaperoning of cancer. Nat. Rev. Cancer 5: 761-772.

### SOURCE

HSP 90 (AC-16) is a mouse monoclonal antibody raised against purified full length native HSP 90 of *Achlya ambisexualis* origin.

# PRODUCT

Each vial contains 200  $\mu g\, lgG_1$  kappa light chain in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

HSP 90 (AC-16) is available conjugated to agarose (sc-101494 AC), 500 µg/ 0.25 ml agarose in 1 ml, for IP; to HRP (sc-101494 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-101494 PE), fluorescein (sc-101494 FITC), Alexa Fluor<sup>®</sup> 488 (sc-101494 AF488), Alexa Fluor<sup>®</sup> 546 (sc-101494 AF546), Alexa Fluor<sup>®</sup> 594 (sc-101494 AF594) or Alexa Fluor<sup>®</sup> 647 (sc-101494 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor<sup>®</sup> 680 (sc-101494 AF680) or Alexa Fluor<sup>®</sup> 790 (sc-101494 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

#### **APPLICATIONS**

HSP 90 (AC-16) is recommended for detection of the constitutive and inducible forms of HSP90 of mouse, rat, human, avian and rabbit origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with the native form of HSP 90 or *E. coli* and yeast HSP 90.

Suitable for use as control antibody for HSP 90 $\alpha/\beta$  siRNA (h): sc-35608, HSP 90 $\alpha/\beta$  siRNA (m): sc-35610, HSP 90 $\alpha/\beta$  shRNA Plasmid (h): sc-35608-SH, HSP 90 $\alpha/\beta$  shRNA Plasmid (m): sc-35610-SH, HSP 90 $\alpha/\beta$  shRNA (h) Lentiviral Particles: sc-35608-V, HSP 90 $\alpha/\beta$  shRNA (m) Lentiviral Particles: sc-35610-V.

#### Positive Controls: HeLa whole cell lysate: sc-2200.

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### SELECT PRODUCT CITATIONS

- 1. Wang, S.A., et al. 2009. Heat shock protein 90 is important for Sp1 stability during mitosis. J. Mol. Biol. 387: 1106-1119.
- Woischnik, M., et al. 2010. A non-BRICHOS surfactant protein c mutation disrupts epithelial cell function and intercellular signaling. BMC Cell Biol. 11: 88.
- 3. Serrano-Marco, L., et al. 2011. Activation of peroxisome proliferator-activated receptor- $\beta$ /- $\delta$  (PPAR- $\beta$ /- $\delta$ ) ameliorates Insulin signaling and reduces SOCS3 levels by inhibiting STAT3 in interleukin-6-stimulated adipocytes. Diabetes 60: 1990-1999.
- 4. Cecarini, V., et al. 2012. Crosstalk between the ubiquitin-proteasome system and autophagy in a human cellular model of Alzheimer's disease. Biochim. Biophys. Acta 1822: 1741-1751.
- 5. Bisicchia, E., et al. 2013. Activation of type-2 cannabinoid receptor inhibits neuroprotective and antiinflammatory actions of glucocorticoid receptor  $\alpha$ : when one is better than two. Cell. Mol. Life Sci. 70: 2191-2204.
- Rajagopal, A., et al. 2015. Proteome of the Insulin-secreting Min6 cell porosome complex: involvement of HSP 90 in its assembly and function. J. Proteomics 114: 83-92.
- 7. Jager, J., et al. 2016. The nuclear receptor Rev-erbα regulates adipose tissue-specific FGF21 signaling. J. Biol. Chem. 291: 10867-10875.
- 8. Yan, M., et al. 2017. Regulation of iNOS-derived Ros generation by HSP 90 and Cav-1 in porcine reproductive and respiratory syndrome virus-infected swine lung injury. Inflammation 40: 1236-1244.
- 9. Karatug Kacar, A. and Bolkent, S. 2018. Necrotic cell death occur via JNK pathway with the activity of transcription factor c-Jun by 4-MC in INS-1 cell line. J. Cell. Biochem. 119: 2048-2060.
- Ullah, M., et al. 2020. HSP 70-mediated NLRP3 inflammasome suppression underlies reversal of acute kidney injury following extracellular vesicle and focused ultrasound combination therapy. Int. J. Mol. Sci. 21: 4085.
- 11. Chen, L., et al. 2020. Mild microwave ablation combined with HSP 90 and TGF- $\beta$ 1 inhibitors enhances the therapeutic effect on osteosarcoma. Mol. Med. Rep. 22: 906-914.
- Öztürk, E., et al. 2020. Thymoquinone is a protective agent that reduces the negative effects of doxorubicin in rat testis. Hum. Exp. Toxicol. 39: 1364-1373.
- Choi, Y.M., et al. 2020. A telomerase-derived peptide exerts an antihepatitis B virus effect via mitochondrial DNA stress-dependent type I interferon production. Front. Immunol. 11: 652.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA