



P23 (JJ12): sc-101497

BACKGROUND

P23, also known as PTGES3 (prostaglandin E synthase 3) or TEBP (telomerase-binding protein p23), is a ubiquitously expressed protein that functions as a cochaperone and plays an important role in signal transduction. One of several proteins in the HSP 90-based molecular chaperone complex, P23 promotes the breakdown of transcriptional regulatory complexes by disrupting receptor-mediated transcriptional activation. P23 acts in a hormone-dependent manner to chaperone estrogen receptor α (ER α), a steroid complex, to its mature form and to regulate the expression of ER α -related genes. Localized to the cytoplasm, P23 interacts with the glucocorticoid receptor (GR) and, through disassembly of the GR transcription machinery, is thought to inhibit GR-dependent transcription. The involvement of P23 in various steroid receptor-mediated pathways suggests close involvement in signal transduction and regulation of cellular processes. Upregulation of P23 is implicated in the invasion and metastasis of various cancers.

REFERENCES

- Muñoz, M.J., et al. 1999. The identification of Wos2, a P23 homologue that interacts with Wee 1 and Cdc2 in the mitotic control of fission yeasts. *Genetics* 153: 1561-1572.
- Freeman, B.C. and Yamamoto, K.R. 2002. Disassembly of transcriptional regulatory complexes by molecular chaperones. *Science* 296: 2232-2235.
- Felts, S.J. and Toft, D.O. 2003. P23, a simple protein with complex activities. *Cell Stress Chaperones* 8: 108-113.
- Tanioka, T., et al. 2003. Regulation of cytosolic prostaglandin E2 synthase by 90 kDa heat shock protein. *Biochem. Biophys. Res. Commun.* 303: 1018-1023.
- Gausdal, G., et al. 2004. Caspase-dependent, geldanamycin-enhanced cleavage of cochaperone P23 in leukemic apoptosis. *Leukemia* 18: 1989-1996.
- Møllerup, J. and Berchtold, M.W. 2005. The cochaperone P23 is degraded by caspases and the proteasome during apoptosis. *FEBS Lett.* 579: 4187-4192.
- Picard, D. 2006. Intracellular dynamics of the HSP 90 co-chaperone P23 is dictated by HSP 90. *Exp. Cell Res.* 312: 198-204.
- Oxelmark, E., et al. 2006. The cochaperone P23 differentially regulates estrogen receptor target genes and promotes tumor cell adhesion and invasion. *Mol. Cell. Biol.* 26: 5205-5213.
- Grad, I., et al. 2006. The HSP 90 co-chaperone p23 is essential for perinatal survival. *Mol. Cell. Biol.* 26: 8976-8983.

CHROMOSOMAL LOCATION

Genetic locus: PTGES3 (human) mapping to 12q13.3; Ptges3 (mouse) mapping to 10 D3.

SOURCE

P23 (JJ12) is a mouse monoclonal antibody raised against P23 of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

P23 (JJ12) is recommended for detection of P23 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for P23 siRNA (h): sc-62741, P23 siRNA (m): sc-62742, P23 shRNA Plasmid (h): sc-62741-SH, P23 shRNA Plasmid (m): sc-62742-SH, P23 shRNA (h) Lentiviral Particles: sc-62741-V and P23 shRNA (m) Lentiviral Particles: sc-62742-V.

Molecular Weight of P23: 23 kDa.

Positive Controls: A549 cell lysate: sc-2413, HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.