

# nucleoporin p62 (2A11): sc-101543

## BACKGROUND

Protein transport across the nucleus is a selective, multi-step process involving several cytoplasmic factors. Proteins must be recognized as import substrates, dock at the nuclear pore complex and translocate across the nuclear envelope in an ATP-dependent fashion. Two cytosolic factors centrally involved in the recognition and docking process are the karyopherin  $\alpha$  and karyopherin  $\beta$  proteins. The karyopherin holoenzyme is a heterodimer of  $\alpha$  and  $\beta$  subunits. Karyopherin  $\alpha$  functions in the recognition and targeting of substrates destined for nuclear import, while karyopherin  $\beta$  serves as an adaptor, tethering the karyopherin  $\alpha$  substrate complex to docking proteins (termed nucleoporins) on the nuclear envelope. p62 glycoprotein is one such nucleoporin, and is not only involved in the nuclear import of proteins, but also the export of nascent mRNA strands. An additional protein, NTF2 (nuclear transport factor 2), interacts with nucleoporin p62 as a homodimer and may be an obligate component of functional p62.

## REFERENCES

- Moroianu, J., et al. 1995. Previously identified protein of uncertain function is karyopherin  $\alpha$  and together with karyopherin  $\beta$  docks import substrate at nuclear pore complexes. *Proc. Natl. Acad. Sci. USA* 92: 2008-2011.
- Moroianu, J., et al. 1995. Protein export from the nucleus requires the GTPase Ran and GTP hydrolysis. *Proc. Natl. Acad. Sci. USA* 92: 4318-4322.
- Buss, F., et al. 1995. Macromolecular interactions in the nucleoporin p62 complex of rat nuclear pores: binding of nucleoporin p54 to the rod domain of p62. *J. Cell Biol.* 128: 251-261.
- Dargemont, C., et al. 1995. Direct interaction of nucleoporin p62 with mRNA during its export from the nucleus. *J. Cell Sci.* 108: 257-263.
- Paschal, B.M., et al. 1995. Identification of NTF2, a cytosolic factor for nuclear import that interacts with nuclear pore complex protein p62. *J. Cell Biol.* 129: 925-937.
- Lounsbury, K.M., et al. 1996. Ran binding domains promote the interaction of Ran with p97/ $\beta$  karyopherin, linking the docking and translocation steps of nuclear import. *J. Biol. Chem.* 271: 2357-2360.
- Moroianu, J., et al. 1996. The binding site of karyopherin  $\alpha$  for karyopherin  $\beta$  overlaps with a nuclear localization sequence. *Proc. Natl. Acad. Sci. USA* 93: 6572-6576.
- Moroianu, J., et al. 1996. Nuclear protein import: Ran-GTP dissociates the karyopherin  $\alpha$ / $\beta$  heterodimer by displacing  $\alpha$  from an overlapping binding site on  $\beta$ . *Proc. Natl. Acad. Sci. USA* 93: 7059-7062.

## CHROMOSOMAL LOCATION

Genetic locus: NUP62 (human) mapping to 19q13.33.

## SOURCE

nucleoporin p62 (2A11) is a rat monoclonal antibody raised against amino acids 1-300 of recombinant nucleoporin p62 of human origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS


nucleoporin p62 (2A11) is recommended for detection of amino acids 1-179 of nucleoporin p62 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for nucleoporin p62 siRNA (h): sc-36107, nucleoporin p62 shRNA Plasmid (h): sc-36107-SH and nucleoporin p62 shRNA (h) Lentiviral Particles: sc-36107-V.


Molecular Weight of nucleoporin p62: 62 kDa.

Positive Controls: nucleoporin p62 (h4): 293T Lysate: sc-158783, HeLa whole cell lysate: sc-2200 or PC-3 cell lysate: sc-2220.

## DATA



nucleoporin p62 (2A11): sc-101543. Western blot analysis of nucleoporin p62 expression in HEK293 (A), BJAB (B), K-562 (C), MOLT-4 (D) and Raji (E) whole cell lysates.



nucleoporin p62 (2A11): sc-101543. Western blot analysis of nucleoporin p62 expression in non-transfected 293T: sc-117752 (A), human nucleoporin p62 transfected 293T: sc-158783 (B), PC-3 (C) and HeLa (D) whole cell lysates.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.