

# Nup153 (R4C8): sc-101545

## BACKGROUND

Nuclear pore complexes (NPCs) are the channels for the bi-directional movement of macromolecules between the nucleus and cytoplasm, and contain more than 100 different subunits. Many of them belong to a family called nucleoporins, which are characterized by the presence of O-linked N-acetylglucosamine moieties and a distinctive pentapeptide repeat (XFXFG). Nup153 is a peripheral NPC component that is implicated in protein and RNP transport and in the interaction of NPCs with the nuclear lamina. Nup153 contains a unique N-terminal region, a central domain consisting of four to five zinc fingers and a C-terminal region containing about 30 irregularly spaced FXFG repeats. Nup153 is cleaved by caspases during apoptosis. Nup153 interacts with TAP, which is essential for mRNA export and associates with chromatin towards the end of anaphase, in parallel with the inner nuclear membrane protein, LAP2. Nup153 is involved in NPC assembly, in anchoring NPCs within the nuclear envelope and in mediating specific nuclear import events.

## REFERENCES

- McMorrow, I., Bastos, R., Horton, H. and Burke, B. 1994. Sequence analysis of cDNA encoding a human nuclear pore complex protein, hNup152. *Biochim. Biophys. Acta* 1217: 219-223.
- Bastos, R., Lin, A., Enarson, M. and Burke, B. 1996. Targeting and function in mRNA export of nuclear pore complex protein Nup153. *J. Cell Biol.* 134: 1141-1156.
- Bodoor, K., Shaikh, S., Salina, D., Raharjo, W.H., Bastos, R., Lohka, M. and Burke, B. 1999. Sequential recruitment of NPC proteins to the nuclear periphery at the end of mitosis. *J. Cell Sci.* 112: 2253-2264.
- Tan, W., Zolotukhin, A.S., Bear, J., Patenaude, D.J. and Felber, B.K. 2000. The mRNA export in *Caenorhabditis elegans* is mediated by CeNXF-1, an ortholog of human TAP/NXF and *Saccharomyces cerevisiae* Mex67p. *RNA* 6: 1762-1772.

## CHROMOSOMAL LOCATION

Genetic locus: NUP153 (human) mapping to 6p22.3; Nup153 (mouse) mapping to 13 A5.

## SOURCE

Nup153 (R4C8) is a rat monoclonal antibody raised against a recombinant protein corresponding to amino acids 610-1191 of Nup153 of rat origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2a</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS


Nup153 (R4C8) is recommended for detection of Nup153 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Nup153 siRNA (h): sc-41279, Nup153 siRNA (m): sc-41280, Nup153 shRNA Plasmid (h): sc-41279-SH, Nup153 shRNA Plasmid (m): sc-41280-SH, Nup153 shRNA (h) Lentiviral Particles: sc-41279-V and Nup153 shRNA (m) Lentiviral Particles: sc-41280-V.


Molecular Weight of Nup153: 150-180 kDa.

Positive Controls: Nup153 (m): 293T Lysate: sc-122173, RAW 264.7 whole cell lysate: sc-2211 or LADMAC whole cell lysate: sc-364189.

## DATA



Nup153 (R4C8): sc-101545. Western blot analysis of Nup153 expression in LADMAC (A), RAW 264.7 (B) and M1 (C) whole cell lysates.



Nup153 (R4C8): sc-101545. Western blot analysis of Nup153 expression in non-transfected 293T: sc-117752 (A), mouse Nup153 transfected 293T: sc-122173 (B) and LADMAC (C) whole cell lysates.

## SELECT PRODUCT CITATIONS

- Kim, S.Y., Ryu, S.J., Ahn, H.J., Choi, H.R., Kang, H.T. and Park, S.C. 2010. Senescence-related functional nuclear barrier by down-regulation of nucleo-cytoplasmic trafficking gene expression. *Biochem. Biophys. Res. Commun.* 391: 28-32.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.