

TFIID (1TBP22): sc-101550

BACKGROUND

In eukaryotic systems, initiation of transcription from protein-coding genes is a complex process requiring RNA polymerase II and broad families of auxiliary transcription factors. Such factors can be divided into two major functional classes: the basal factors that are required for transcription of all Pol II genes, including TFIIA, TFIIIB, TFIID, TFIIIE, TFIIF and TFIIH; and sequence-specific factors that regulate gene expression. The basal transcription factors and Pol II form a specific multiprotein complex near the transcription start site by interacting with core promoter elements such as the TATA box generally located 25-30 base pairs upstream of the transcription start site. Binding of TFIID to the TATA element initiates assembly of the other factors into a pre-initiation complex. The TATA-binding subunit of TFIID (designated TFIIDt or TBP) from higher eukaryotes contains a highly conserved 180 amino acid C-terminal domain.

REFERENCES

- Maldonado, E., et al. 1990. Factors involved in specific transcription by mammalian RNA polymerase II: role of transcription factors IIA, IID, and IIB during formation of a transcription-competent complex. *Mol. Cell. Biol.* 10: 6335-6347.
- Peterson, M.G., et al. 1991. Structure and functional properties of human general transcription factor IIE. *Nature* 354: 369-373.
- Lee, D.K., et al. 1992. TFIIA induces conformational changes in TFIID via interactions with the basic repeat. *Mol. Cell. Biol.* 12: 5189-5196.
- Takada, R., et al. 1992. Identification of human TFIID components and direct interaction between a 250 kDa polypeptide and the TATA box-binding protein (TFIIDt). *Proc. Natl. Acad. Sci. USA* 89: 11809-11813.
- Huisinga, K.L., et al. 2007. A TATA binding protein regulatory network that governs transcription complex assembly. *Genome Biol.* 8: R46.
- Romier, C., et al. 2007. Crystal structure, biochemical and genetic characterization of yeast and *E. cuniculi* TAF(II)5 N-terminal domain: implications for TFIID assembly. *J. Mol. Biol.* 368: 1292-1306.
- Demeny, M.A., et al. 2007. Identification of a small TAF complex and its role in the assembly of TAF-containing complexes. *PLoS ONE* 2: 316.
- Bhattacharya, S., et al. 2007. Structural analysis and dimerization potential of the human TAF5 subunit of TFIID. *Proc. Natl. Acad. Sci. USA* 104: 1189-1194.

CHROMOSOMAL LOCATION

Genetic locus: TBP (human) mapping to 6q27; Tbp (mouse) mapping to 17 A2.

SOURCE

TFIID (1TBP22) is a mouse monoclonal antibody raised against amino acids 55-99 of TFIID of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

TFIID (1TBP22) is recommended for detection of TFIID of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for TFIID siRNA (h): sc-29503, TFIID siRNA (m): sc-36648, TFIID shRNA Plasmid (h): sc-29503-SH, TFIID shRNA Plasmid (m): sc-36648-SH, TFIID shRNA (h) Lentiviral Particles: sc-29503-V and TFIID shRNA (m) Lentiviral Particles: sc-36648-V.


Molecular Weight of TFIID: 36 kDa.

Positive Controls: TFIID (m): 293T Lysate: sc-124002, TFIID (h): 293T Lysate: sc-111938 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



TFIID (1TBP22): sc-101550. Western blot analysis of TFIID expression in non-transfected 293T: sc-117752 (A), human TFIID transfected 293T: sc-111938 (B) and HeLa (C) whole cell lysates.



TFIID (1TBP22): sc-101550. Western blot analysis of TFIID expression in non-transfected: sc-117752 (A) and mouse TFIID transfected: sc-124002 (B) 293T whole cell lysates.

STORAGE


Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **TFIID (TBP) (58C9): sc-421** for TFIID antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.