

VRK1 (1F6): sc-101554

BACKGROUND

Human vaccinia-related kinases 1 and 2 (VRK1/2) are NLS-containing, serine/threonine poxvirus-related kinases that are similar to casein kinase I family members. These VRK kinases phosphorylate transcription factors related to stress responses, such as p53. As an upstream regulator of p53, VRK1 is capable of phosphorylating phospho-tyrosine, casein, histone 2b and myelin basic protein. VRK1 colocalizes with ATF-2 in the nucleus and can form a stable complex. VRK1 phosphorylates ATF-2 mainly on Thr-73, stabilizing the ATF-2 protein and increasing its intracellular level. VRK1 phosphorylates human p53 in Thr-18 and disrupts p53-MDM2 interaction *in vitro*. VRK1 phosphorylates c-Jun in Ser-63 and Ser-73 *in vitro* (the same residues targeted by the N-terminal kinase of c-Jun (JNK)), and activates c-Jun dependent transcription.

REFERENCES

- Hunter, T. 1995. Protein kinases and phosphatases: the yin and yang of protein phosphorylation and signaling. *Cell* 80: 225-236.
- Nezu, J., et al. 1997. Identification of two novel human putative serine/threonine kinases, VRK1 and VRK2, with structural similarity to Vaccinia Virus B1R kinase. *Genomics* 45: 327-331.
- Lopez-Borges, S. and Lazo, P.A. 2000. The human vaccinia-related kinase 1 (VRK1) phosphorylates Threonine-18 within the MDM2 binding site of the p53 tumour suppressor protein. *Oncogene* 19: 3656-3664.
- Nichols, R.J., et al. 2004. Characterization of three paralogous members of the mammalian vaccinia related kinase family. *J. Biol. Chem.* 279: 7934-7946.
- Boyle, K.A., et al. 2004. Members of a novel family of mammalian protein kinases complement the DNA-negative phenotype of a Vaccinia Virus *ts* mutant defective in the B1 kinase. *J. Virol.* 78: 1992-2005.
- Sevilla, A., et al. 2004. Human vaccinia-related kinase 1 (VRK1) activates the ATF-2 transcriptional activity by novel phosphorylation on Thr-73 and Ser-62 and cooperates with JNK. *J. Biol. Chem.* 279: 27458-27465.
- Sevilla, A., et al. 2004. c-Jun phosphorylation by the human vaccinia-related kinase 1 (VRK1) and its cooperation with the N-terminal kinase of c-Jun (JNK). *Oncogene* 23: 8950-8958.

CHROMOSOMAL LOCATION

Genetic locus: VRK1 (human) mapping to 14q32.2; Vrk1 (mouse) mapping to 12 F1.

SOURCE

VRK1 (1F6) is a mouse monoclonal antibody raised against recombinant VRK1 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

VRK1 (1F6) is recommended for detection of VRK1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for VRK1 siRNA (h): sc-106702, VRK1 siRNA (m): sc-155227, VRK1 shRNA Plasmid (h): sc-106702-SH, VRK1 shRNA Plasmid (m): sc-155227-SH, VRK1 shRNA (h) Lentiviral Particles: sc-106702-V and VRK1 shRNA (m) Lentiviral Particles: sc-155227-V.


Molecular Weight of VRK1: 47 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, VRK1 (m2): 293T Lysate: sc-124593 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



VRK1 (1F6): sc-101554. Western blot analysis of VRK1 expression in HeLa (A), HL-60 (B) and HCT-116 (C) whole cell lysates.



VRK1 (1F6): sc-101554. Western blot analysis of VRK1 expression in non-transfected 293T: sc-117752 (A), mouse VRK1 transfected 293T: sc-124593 (B) and HeLa (C) whole cell lysates.

SELECT PRODUCT CITATIONS

- Valbuena, A., et al. 2008. Human VRK1 is an early response gene and its loss causes a block in cell cycle progression. *PLoS ONE* 3: e1642.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.