

α -defensin 6 siRNA (h): sc-105018

BACKGROUND

Human neutrophil α -defensins (also designated HNPs) are small, cationic, cysteine-rich antimicrobial proteins that play important roles in innate immunity against infectious microbes such as bacteria, fungi and enveloped viruses. α -defensins are synthesized as inactive precursors and are activated by proteolytic cleavage by MMP-7. Paneth cells in small intestinal crypts secrete the α -defensins, which are also termed cryptidins in mice. α -defensins 5 and 6 probably contribute to innate defense of the GI mucosal surface by protecting against microbial invasion in states of intestinal inflammation.

REFERENCES

1. Ouellette, A.J., et al. 1999. Peptide localization and gene structure of cryptidin 4, a differentially expressed mouse paneth cell α -defensin. *Infect. Immun.* 67: 6643-6651.
2. Ayabe, T., et al. 2002. Activation of Paneth cell α -defensins in mouse small intestine. *J. Biol. Chem.* 277: 5219-5228.
3. Wu, Z., et al. 2003. From pro defensins to defensins: synthesis and characterization of human neutrophil pro α -defensin-1 and its mature domain. *J. Pept. Res.* 62: 53-62.
4. Cunliffe, R.N. 2003. α -defensins in the gastrointestinal tract. *Mol. Immunol.* 40: 463-467.
5. Maemoto, A., et al. 2004. Functional analysis of the α -defensin disulfide array in mouse cryptidin-4. *J. Biol. Chem.* 279: 44188-44196.
6. Nam, M.J., et al. 2004. Identification of defensin α 6 as a potential biomarker in colon adenocarcinoma. *J. Biol. Chem.* 280: 8260-8265.
7. Wu, Z., et al. 2005. Human neutrophil α -defensin 4 inhibits HIV-1 infection *in vitro*. *FEBS Lett.* 579: 162-166.

CHROMOSOMAL LOCATION

Genetic locus: DEFA6 (human) mapping to 8p23.1.

PRODUCT

α -defensin 6 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see α -defensin 6 shRNA Plasmid (h): sc-105018-SH and α -defensin 6 shRNA (h) Lentiviral Particles: sc-105018-V as alternate gene silencing products.

For independent verification of α -defensin 6 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105018A and sc-105018B.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

α -defensin 6 siRNA (h) is recommended for the inhibition of α -defensin 6 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.