

AChR α 10 siRNA (h): sc-105030

BACKGROUND

Members of the ligand-gated ion channel receptor family are characterized by their fast transmitting response to neurotransmitters. Two important members of this family are the nicotinic acetylcholine and glutamate receptors, both of which are composed of five homologous subunits forming a transmembrane aqueous pore. These transmembrane receptors change conformation in response to their cognate neurotransmitter. Nicotinic acetylcholine receptors (AChRs) are found at the postsynaptic membrane of the neuromuscular junction and bind acetylcholine molecules, allowing ions to move through the pore. Glutamate receptors are found in the postsynaptic membrane of cells in the central nervous system. The activity that is generated at the synapse by the binding of acetylcholine is terminated by acetylcholinesterase, an enzyme that rapidly hydrolyzes acetylcholine. AChR α 10, also known as CHRNA10, is a 450 amino acid multi-pass membrane protein expressed in inner-ear tissue, tonsil, immortalized B-cells, cultured T-cells and peripheral blood lymphocytes. AChR α 10 forms a hetero-oligomeric channels in conjunction with AChR α 9 and is considered an ionotropic receptor with a probable role in the modulation of auditory stimuli.

REFERENCES

1. Alkondon, M., et al. 1988. Acetylcholinesterase reactivators modify the functional properties of the nicotinic acetylcholine receptor ion channel. *J. Pharmacol. Exp. Ther.* 245: 543-556.
2. Betz, H. 1990. Ligand-gated ion channels in the brain: the amino acid receptor superfamily. *Neuron* 5: 383-392.
3. Baenziger, J.E., et al. 1992. Probing conformational changes in the nicotinic acetylcholine receptor by Fourier transform infrared difference spectroscopy. *Biophys. J.* 62: 64-66.
4. Daw, N.W., et al. 1993. The role of NMDA receptors in information processing. *Annu. Rev. Neurosci.* 16: 207-222.
5. Stevens, C.F. 1993. Quantal release of neurotransmitter and long-term potentiation. *Cell* 72: 55-63.

CHROMOSOMAL LOCATION

Genetic locus: CHRNA10 (human) mapping to 11p15.4.

PRODUCT

AChR α 10 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AChR α 10 shRNA Plasmid (h): sc-105030-SH and AChR α 10 shRNA (h) Lentiviral Particles: sc-105030-V as alternate gene silencing products.

For independent verification of AChR α 10 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105030A, sc-105030B and sc-105030C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AChR α 10 siRNA (h) is recommended for the inhibition of AChR α 10 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AChR α 10 gene expression knockdown using RT-PCR Primer: AChR α 10 (h)-PR: sc-105030-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.