

AP-3 σ 2 siRNA (h): sc-105076

BACKGROUND

AP-3 σ 2, also known as AP3S2 (adaptor-related protein complex 3, σ 2 subunit), AP-3 complex σ -3B subunit or clathrin-associated/assembly/adaptor protein, small 4, 22-kD, is a 193 amino acid protein that belongs to the adaptor complexes small subunit family and is ubiquitously expressed. Encoded by a gene that maps to human chromosome 15q26.1, AP-3 σ 2 interacts with AGAP1 and may play a role in carotid plaque traits. AP-3 σ 2 is part of the adapter protein complex 3 (AP-3), which is a heterotetramer that consists of two large adaptins (AP-3 δ and AP-3 β or β -NAP), one medium adaptin (AP-3 μ or AP-3 μ 2) and one small adaptin (σ -type subunit APS1 or AP-3 σ 2). Associated with the Golgi region and peripheral structures, AP-3 σ 2 assists in vesicle budding from the Golgi membrane and may be highly engaged in lysosome trafficking.

REFERENCES

- Dell'Angelica, E.C., Ohno, H., Ooi, C.E., Rabinovich, E., Roche, K.W. and Bonifacino, J.S. 1997. AP-3: an adaptor-like protein complex with ubiquitous expression. *EMBO J.* 16: 917-928.
- Lefrancois, S., Janvier, K., Boehm, M., Ooi, C.E. and Bonifacino, J.S. 2004. An ear-core interaction regulates the recruitment of the AP-3 complex to membranes. *Dev. Cell* 7: 619-625.
- Belbin, T.J., Singh, B., Smith, R.V., Socci, N.D., Wreesmann, V.B., Sanchez-Carbayo, M., Masterson, J., Patel, S., Cordon-Cardo, C., Prystowsky, M.B. and Childs, G. 2005. Molecular profiling of tumor progression in head and neck cancer. *Arch. Otolaryngol. Head Neck Surg.* 131: 10-18.
- Chittenden, T.W., Sherman, J.A., Xiong, F., Hall, A.E., Lanahan, A.A., Taylor, J.M., Duan, H., Pearlman, J.D., Moore, J.H., Schwartz, S.M. and Simons, M. 2006. Transcriptional profiling in coronary artery disease: indications for novel markers of coronary collateralization. *Circulation* 114: 1811-1820.
- Chen, C., Zou, X., Ji, C., Zhao, S., Lv, L., Gu, S., Xie, Y. and Mao, Y. 2007. Characterization of AP3B2_v2, a novel splice variant of human AP3B2. *DNA Seq.* 18: 165-168.
- Li, Y., Chang, M., Abar, O., Garcia, V., Rowland, C., Catanese, J., Ross, D., Broder, S., Shiffman, M., Cheung, R., Wright, T., Friedman, S.L. and Sninsky, J. 2009. Multiple variants in Toll-like receptor 4 gene modulate risk of liver fibrosis in Caucasians with chronic hepatitis C infection. *J. Hepatol.* 51: 750-757.
- Mead, C.L., Kuzyk, M.A., Moradian, A., Wilson, G.M., Holt, R.A. and Morin, G.B. 2010. Cytosolic protein interactions of the schizophrenia susceptibility gene dysbindin. *J. Neurochem.* 113: 1491-1503.
- Dong, C., Beecham, A., Slifer, S., Wang, L., Blanton, S.H., Wright, C.B., Rundek, T. and Sacco, R.L. 2010. Genomewide linkage and peakwise association analyses of carotid plaque in caribbean hispanics. *Stroke* 41: 2750-2756.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: AP3S2 (human) mapping to 15q26.1.

PRODUCT

AP-3 σ 2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see AP-3 σ 2 shRNA Plasmid (h): sc-105076-SH and AP-3 σ 2 shRNA (h) Lentiviral Particles: sc-105076-V as alternate gene silencing products.

For independent verification of AP-3 σ 2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-105076A, sc-105076B and sc-105076C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

AP-3 σ 2 siRNA (h) is recommended for the inhibition of AP-3 σ 2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor AP-3 σ 2 gene expression knockdown using RT-PCR Primer: AP-3 σ 2 (h)-PR: sc-105076-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.